

Clinical Reasoning: A 26-Year-Old Woman With Recurrent Pain, Weakness, and Atrophy in Bilateral Upper Limbs During Pregnancy and Puerperium

Tian-fang Zeng, MD,* Xin Ding, MD,* Dan Yang, MD, Chang-Chuan Wu, MD, Dan-Dan Xie, MD, and Wen-Min Zhang, MD

Neurology® 2023;100:631-637. doi:10.1212/WNL.0000000000201705

Correspondence

Dr. Ding
dingxin105@163.com

Abstract

We present the case of a 26-year-old woman with recurrent episodes of severe pain, weakness, and atrophy in her bilateral upper extremities during pregnancy and puerperium. She reported 2 similar episodes at ages 5 and 10 years, after which she fully recovered. On examination, we observed significant atrophy in her bilateral upper extremity muscles with decreased strength. Needle electromyography (EMG) revealed neurogenic damage in her bilateral upper limbs. The patient's clinical manifestations and auxiliary examination suggested a brachial plexopathy. Metabolic and immune factors that may occur during pregnancy and puerperium were evaluated. We also screened for paraneoplastic, neoplastic, and genetic factors. Finally, a hereditary form of disease was considered. This case emphasizes the importance of early diagnosis and avoidance of triggers.

*T.-f. Zeng and X. Ding are co-first authors.

From the Department of Neurology (T.-f.Z., X.D., D.Y., C.-C.W., D.-D.X.), Cheng Du Second People's Hospital; and Chengdu Medical College (W.-M.Z.), China.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Section 1

A 26-year-old woman developed tingling in her right upper limb during the second month of pregnancy that progressed to her left upper limb within several days. The pain, which first appeared at night, worsened during cold weather and improved when it was hot. Two weeks later, she developed proximal weakness in her right more than left arm. She also noticed progressive difficulty lifting her arms, using chopsticks, and buttoning clothes. There was no history of antecedent trauma, infection, or vaccination. The patient did not take any medication and was referred for rehabilitation therapy. The pain gradually resolved 3 months later, and the weakness in both arms slowly improved. She was eventually able to manage her daily activities without assistance, but had mild residual weakness.

One week after giving birth, the patient experienced another attack. She developed severe pain in the left shoulder and arm, numbness in the left hand, and could not raise her left arm. After a few days, she developed weakness and pain in both upper limbs. No neck pain, bowel/bladder dysfunction, or muscle fasciculations were


reported. She denied weakness or sensory change in her lower limbs. She reported having 2 similar episodes at ages 5 and 10 years, after which she fully recovered. Regarding family history, her mother also reported a similar single episode that started with muscle weakness in the right upper extremity 1 month after parturition, from which she recovered 8 months later.

On examination, mental status was normal and cranial nerves were intact. Strength in the upper extremities was decreased (Figure, A). The patient had normal strength in the lower extremities. Tone was decreased in the arms and normal in the legs. Significant atrophy of bilateral infraspinatus, teres minor, and deltoid muscles were observed in addition to decreased sensation to pinprick along the radial aspect of the forearms. No fasciculations were noted. Deep tendon reflexes (DTRs) were 1+ in the upper limbs and 2+ in the lower limbs. No pathologic reflexes were found. The patient had a similar facial appearance as her mother because both had hypotelorism (measured pupil-to-pupil with a ruler) (Figure, B).

Question for Consideration:

1. Where would you localize the lesion?

Figure Clinical and Gene Findings



(A) Strength in the upper extremities (Medical Research Council); (B) Dysmorphic features; (C) A heterozygous single nucleotide change c.316C > T in exon 3 of the *SEPT9* gene, resulting in the amino acid change p.R106W; (D) Pedigree: Female individuals are shown as circles, male individuals as squares. Filled symbols represent affected individuals with an exon 3 missense variation of the *SEPT9* gene. Open symbols indicate nonaffected individuals and a slanting arrow the index patient. III-1 is without attack so far.

GO TO SECTION 2

Section 2

The severe pain, numbness, and weakness with focal muscle atrophy in the bilateral upper limbs and depressed reflexes suggest involvement of the peripheral nervous system. The CNS is less likely involved in the absence of upper motor neuron signs. Therefore, we considered a multilevel radicular process involving multiple nerve roots (C5-T1), brachial plexopathy, or neuropathy (radial, median, axillary, supra-scapular, and musculocutaneous nerves). The patient also experienced significant pain during each attack. The differential diagnosis based on localization might involve the following:

1. Radiculopathy, which can be caused by trauma or various inflammatory or neoplastic disorders but does not typically involve the bilateral upper extremities simultaneously.
2. Plexopathy, such as Parsonage-Turner syndrome or diabetic radiculoplexus neuropathies, which can be associated with severe pain at the onset of weakness.
3. Multiple mononeuropathies, which can be caused by vasculitis, inflammation, or diabetic neuropathy and result in neuropathic pain.

Question for Consideration:

1. What is the most appropriate next step in this patient's workup?

GO TO SECTION 3

Section 3

Needle electromyography (EMG) and nerve conduction studies (NCS) were performed after the fourth attack. EMG revealed neurogenic damage in the bilateral upper limbs. Sensory nerve conduction studies also showed a low-amplitude right radial nerve sensory nerve action potential (SNAP) and slow sensory conduction velocity of the right median nerve and bilateral radial nerves. The motor nerve conduction testing showed decreased compound muscle action potential (CMAP) amplitudes elicited bilaterally from the axillary, radial, and musculocutaneous nerves and right median nerve (Table). The abovementioned findings revealed dysfunction in the middle and upper trunci of the bilateral brachial plexus.

The patient's clinical manifestations and auxiliary examination suggested a brachial plexus neuropathy; however,

she had no history of recent infection, trauma, vaccination, cancer, toxic exposure, diabetes, or radiation therapy. Laboratory tests, including erythrocyte sedimentation rate, antinuclear antibodies, and antineutrophil cytoplasmic antibodies, were within normal limits. CSF tests were normal. Immunologic testing for antiganglioside and anti-Hu paraneoplastic antibodies showed negative results. Contrast-enhanced MRI of the brachial plexus and cervical spine was normal. Considering the high rate of recurrence, family history, and presence of dysmorphic features, a hereditary disease was suspected.

Questions for Consideration:

1. What hereditary disease could account for patient's symptoms?
2. What testing is required for definitive diagnosis?

Table Nerve Conduction Studies and EMG

| CMAP | Stimulation site | Recording site | Latency, ms | Amplitude (motor mV, sensory μ V) | Conduction velocity, m/s |
|---------------------------------|-----------------------------|----------------------|------------------------------|---------------------------------------|--------------------------|
| R median | Wrist | APB | 3.80 (<4) | 0.7 (>5) | |
| | Elbow | APB | 7.81 | 0.4 (>5) | 54 (>50) |
| L median | Wrist | APB | 2.50 (<4) | 6.4 (>5) | |
| | Elbow | APB | 5.52 | 5.5 (>5) | 73 (>50) |
| R ulnar | Wrist | ADM | 1.88 (<3.8) | 7.1 (>5) | |
| | Below elbow | ADM | 5.00 | 8.2 (>5) | 74 (>50) |
| L ulnar | Wrist | ADM | 2.24 (<3.8) | 11.2 (>5) | |
| | Below elbow | ADM | 5.05 | 9.4 (>5) | 82 (>50) |
| R radial | Forearm | EIP | 4.06 | 0.3 (>8) | — |
| L radial | Forearm | EIP | 4.38 | 0.3 (>8) | — |
| R musculocutaneous nerve | Erb's | Biceps | 5.99 | 1.4 (>6) | — |
| L musculocutaneous nerve | Erb's | Biceps | 4.84 | 1.2 (>6) | — |
| R axillary nerve | Erb's | Deltoid | 3.23 | 1.2 (>6) | — |
| L axillary nerve | Erb's | Deltoid | 4.48 | 3.0 (>6) | — |
| R tibial | Ankle | AH | 2.71 (<5) | 18.0 (>4) | |
| L peroneal | Ankle | EDB | 3.80 (<5) | 6.3 (>3) | |
| SNAP | | | | | |
| R median | Wrist | Digit II | 5.94 | 43.7 (>10) | 26 (>50) |
| L median | Wrist | Digit II | 2.45 | 65.0 (>10) | 65 (>50) |
| R ulnar | Wrist | Digit V | 2.34 | 56.3 (>10) | 58 (>50) |
| L ulnar | Wrist | Digit V | 2.19 | 32.0 (>10) | 60 (>50) |
| R radial | Forearm | Anatomical snuff box | 2.97 | 2.90 (>10) | 45 (>50) |
| L radial | Forearm | Anatomical snuff box | 2.92 | 20.5 (>10) | 45 (>50) |
| EMG | Spontaneous activity | | Motor unit morphology | | |
| | Fibs/PSWs | Fasciculation | Amplitude | Duration | Recruitment |
| L deltoid | 3+ | — | No units | No units | No units |

Continued

Table Nerve Conduction Studies and EMG *(continued)*

| EMG | Spontaneous activity | | Motor unit morphology | | |
|----------------------------|----------------------|---------------|-----------------------|-----------|-------------|
| | Fibs/PSWs | Fasciculation | Amplitude | Duration | Recruitment |
| L biceps | 1+ | — | Normal | Normal | Decreased |
| L EDC | 2+ | — | No units | No units | No units |
| L APB | 1+ | — | No units | No units | No units |
| R deltoid | 2+ | — | Normal | Normal | No units |
| R biceps | 2+ | — | Increased | Increased | Decreased |
| R EDC | 3+ | — | Increased | Increased | No units |
| R APB | 3+ | — | No units | No units | No units |
| L tibialis anterior | — | — | Normal | Normal | Normal |
| R vastus medialis | — | — | Normal | Normal | Normal |

Abbreviations: ADM = abductor digiti minimi; AH = abductor hallucis; APB = abductor pollicis brevis; CMAP = compound muscle action potential; EDB = extensor digitorum brevis; EDC = extensor digitorum communis; EIP = extensor indicis proprius; EMG = Needle electromyography; Fibs/PSWs = fibrillation potentials or positive sharp waves; SNAP = sensory nerve action potential.

Annotation: The values in parentheses are reference normal values.

GO TO SECTION 4

Section 4

When evaluating patients for a genetic disease involving the brachial plexus, hereditary neuralgic amyotrophy (HNA) should be considered. Gene analyses were performed for HNA. We identified a heterozygous single nucleotide change c.316C > T in exon 3 of the *SEPT9* gene (GRCh37/hg19 chr17:75398380), resulting in the amino acid change p.R106W (Figure, C). This variation was also found in the patient's mother and son (Figure, D). The son was 2 months old during genetic testing.

Discussion

Neuralgic amyotrophy (NA) is clinically characterized by the sudden onset of extreme neuropathic pain in the upper limbs, followed by weakness and atrophy of the affected muscles and occasional sensory deficits, with slow recovery over months to years.¹ Neuralgic amyotrophy has an idiopathic form (INA, also called Parsonage-Turner syndrome) and an autosomal dominant hereditary form (HNA). Compared with INA, HNA is much rarer.^{2,3} Single HNA episodes strongly resemble INA. The striking similarities include pain, predilection for the arm plexus, and provocation by the same triggers.⁴ Distinguishing features include family history, earlier age at onset, a higher rate of recurrence, and the presence of dysmorphic features in HNA, which are noteworthy in our case.

The European CMT Consortium developed the first diagnostic guidelines for HNA.⁵ Subsequently, Alfen et al.⁶ supplemented the characteristics of HNA. They proposed core features of HNA, which are as follows: (1) acute, unilateral, or bilateral brachial plexopathy; (2) severe pain precedes onset of weakness by days to few weeks; (3) predominantly motor deficits; (4) number of episodes variable (1–20); and (5) precipitating factors: infections, immunizations, surgery, pregnancy, parturition, unusually strenuous exercise of the affected limb, and exposure to cold. In our study, the patient suffered from attacks during pregnancy and after delivery. The onset of HNA usually occurs in the second or third decade of life, though earlier and later onset is possible. HNA can run 2 distinct courses: a relapsing/remitting course with symptom-free intervals or incomplete recovery, characterized by persistent neurologic deficits after repeated attacks in the same limb.^{6,7}

Several authors have also noted minor dysmorphic features associated with HNA. These include a long, narrow face, small mouth, hypotelorism (close-set eyes), shortened palpebral fissures, epicanthal folds, cleft palate, minor syndactyly, circular skin creases, long nasal bridge, and short stature.^{8,9} Both our patient and her mother have mild dysmorphic features, such as hypotelorism, which can help differentiate HNA from INA.⁵

Regarding genetics, HNA (Online Mendelian Inheritance in Man catalog, OMIM 162100) is an autosomal dominant disorder associated with pathogenic variations in the *SEPT9* gene on chromosome 17q25.3.^{10, 11} To date, 3 point variations

(c.-131G > C, c.262C > T, and c.278C > T) and a genetic founder haplotype have been identified in HNA pedigrees supporting a critical role for the *SEPT9* gene.^{8,12,13} In general, HNA is genetically heterogeneous and has been linked to a variation or duplication in the *SEPT9* gene in only 55% of affected families.¹⁴ One or more unknown genes may also be associated with HNA.³

So far, sequence analysis of *SEPT9* in larger cohorts of HNA families has not identified any additional variations beyond those previously reported.^{11,14} In this study, our patient had an unusual heterozygous variation with typical HNA symptoms, NM_001113491.1p.R106W (c.316C > T), located on exon 3 of the *SEPT9* gene. Genetic analyses suggested a heterozygous variation in the *SEPT9* gene in 3 generations, the patient, her mother, and her son.

Currently, there is no standardized approach for treatment. Some case series suggest that early corticosteroid therapy or i.v. immunoglobulin may benefit patients in the acute phase of NA.¹⁵ This case emphasizes the importance of early diagnosis and avoidance of triggers. HNA should be highly suspected when a patient presents with severe pain followed by weakness and atrophy of the upper extremities. Although we found a gene locus that might be responsible for HNA, the significance of this finding should be verified in subsequent studies.

Study Funding

The authors report no targeted funding.

Disclosure

The authors report no relevant disclosures. Go to Neurology.org/N for full disclosures.

Publication History

Received by *Neurology* April 24, 2022. Accepted in final form November 3, 2022. Submitted and externally peer reviewed. The handling editor was Resident and Fellow Section Editor Whitley Aamodt, MD, MPH.

Appendix Authors

| Name | Location | Contribution |
|--------------------|--|--|
| Tian-fang Zeng, MD | Department of Neurology, Cheng Du Second People's Hospital, Chengdu, China | Drafting/revision of the article for content, including medical writing for content |
| Xin Ding, MD | Department of Neurology, Cheng Du Second People's Hospital, Chengdu, China | Drafting/revision of the article for content, including medical writing for content; study concept or design |
| Dan Yang, MD | Department of Neurology, Cheng Du Second People's Hospital, Chengdu, China | Study concept or design |
| Chang-Chuan Wu, MD | Department of Neurology, Cheng Du Second People's Hospital, Chengdu, China | Major role in the acquisition of data |

Appendix (continued)

| Name | Location | Contribution |
|-------------------|--|------------------------------------|
| Dan-Dan Xie, MD | Department of Neurology, Cheng Du Second People's Hospital, Chengdu, China | Analysis or interpretation of data |
| Wen-Min Zhang, MD | Chengdu Medical College, Chengdu, China | Analysis or interpretation of data |

References

1. van Alfen N, van Eijk JJJ, Ennik T, et al. Incidence of neuralgic amyotrophy (Parsonage Turner syndrome) in a primary care setting—a prospective cohort study. *PLoS One* 2015;10(5), e0128361, 10.1371/journal.pone.0128361
2. Parsonage MJ, Aldren Turner J. Neuralgic amyotrophy; the shoulder-girdle syndrome. *The Lancet* 1948;251(6513), 973-978, 10.1016/s0140-6736(48)90611-4
3. Neubauer K, Boeckelmann D, Koehler U, et al. Hereditary neuralgic amyotrophy in childhood caused by duplication within the SEPT9 gene: a family study. *Cytoskeleton (Hoboken)* 2019;76(1), 131-136, 10.1002/cm.21479
4. Meuleman J, Timmerman V, Van Broeckhoven C, De Jonghe P. Hereditary neuralgic amyotrophy. *Neurogenetics* 2001;3:115-118, 10.1007/s100480100109
5. Kuhlenbäumer G, Stögbauer F, Timmerman V, De Jonghe P. Diagnostic guidelines for hereditary neuralgic amyotrophy or hereditary familial neuritis with brachial plexus predilection. *Neuromuscul Disord.* 2000;10(7), 515-517, 10.1016/s0960-8966(00)00109-7
6. van Alfen N, Hannibal MC, Chance PF, van Engelen BGM. Hereditary neuralgic amyotrophy—retired chapter, for historical reference only. In: Adam MP, Everman DB, Mirzaa GM, et al., eds. *GeneReviews*(®): University of Washington, Seattle Copyright © 1993-2022, University of Washington, Seattle. GeneReviews is a registered trademark of the University of Washington, Seattle. All rights reserved, 1993.
7. van Alfen N, van Engelen BG, Reinders JW, Kremer H, Gabreëls FJ. The natural history of hereditary neuralgic amyotrophy in the Dutch population: two distinct types? *Brain* 2000;123 (4):718-723, 10.1093/brain/123.4.718
8. Laccone F, Hannibal MC, Neesen J, Grisold W, Chance PF, Rehder H. Dysmorphic syndrome of hereditary neuralgic amyotrophy associated with a SEPT9 gene mutation—a family study. *Clin Genet.* 2008;74(3), 279-283, 10.1111/j.1399-0004.2008.01022.x
9. Jeannot PY, Watts GD, Bird TD, Chance PF. Craniofacial and cutaneous findings expand the phenotype of hereditary neuralgic amyotrophy. *Neurology* 2001;57(11), 1963-1968, 10.1212/wnl.57.11.1963
10. Wehnert M, Timmerman V, Spoelders P, Meuleman J, Nelis E, Van Broeckhoven C. Further evidence supporting linkage of hereditary neuralgic amyotrophy to chromosome 17q. *Neurology* 1997;48(6), 1719-1721, 10.1212/wnl.48.6.1719
11. Hannibal MC, Ruzzo EK, Miller LR, et al. SEPT9 gene sequencing analysis reveals recurrent mutations in hereditary neuralgic amyotrophy. *Neurology* 2009;72(20), 1755-1759, 10.1212/wnl.0b013e3181a609e3
12. Kuhlenbäumer G, Hannibal MC, Nelis E, et al. Mutations in SEPT9 cause hereditary neuralgic amyotrophy. *Nat Genet.* 2005;37(10), 1044-1046, 10.1038/ng1649
13. Ueda M, Kawamura N, Tateishi T, et al. Phenotypic spectrum of hereditary neuralgic amyotrophy caused by the SEPT9 R88W mutation. *J Neurol Neurosurg Psychiatry* 2010;81(1), 94-96, 10.1136/jnnp.2008.168260
14. Collie AMB, Landsverk ML, Ruzzo E, et al. Non-recurrent SEPT9 duplications cause hereditary neuralgic amyotrophy. *J Med Genet.* 2010;47(9), 601-607, 10.1136/jmg.2009.072348
15. IJspeert J, Janssen RMJ, van Alfen N. Neuralgic amyotrophy. *Curr Opin Neurol.* 2021; 34(5), 605-612, 10.1097/wco.0000000000000968

Share Your Artistic Expressions in Neurology ‘Visions’

AAN members are urged to submit medically or scientifically related artistic images, such as photographs, photomicrographs, and paintings, to the “Visions” section of *Neurology*®. These images are creative in nature, rather than the medically instructive images published in the NeuroImages section. The image or series of up to six images may be black and white or color and must fit into one published journal page. Accompanying description should be 100 words or less; the title should be a maximum of 140 characters including spaces and punctuation.

Please access the Author Center at [Npub.org/authors](https://www.npub.org/authors) for full submission information.

NEW EPISODE

Neurology®
PODCAST



Listen and Learn on the Go With the Neurology® Podcast!

Access from your smart speaker, car, or just about anywhere. Download, subscribe, and rate/review the *Neurology* Podcast wherever you enjoy your podcasts. Head to [Neurology.libsyn.com/website](https://www.neurology.com/libsyn.com/website) for the full list of past episodes.

Neurology®

Clinical Reasoning: A 26-Year-Old Woman With Recurrent Pain, Weakness, and Atrophy in Bilateral Upper Limbs During Pregnancy and Puerperium

Tian-fang Zeng, Xin Ding, Dan Yang, et al.

Neurology 2023;100:631-637 Published Online before print December 15, 2022

DOI 10.1212/WNL.0000000000201705

This information is current as of December 15, 2022

| | |
|---|---|
| Updated Information & Services | including high resolution figures, can be found at: http://n.neurology.org/content/100/13/631.full |
| References | This article cites 14 articles, 5 of which you can access for free at: http://n.neurology.org/content/100/13/631.full#ref-list-1 |
| Subspecialty Collections | This article, along with others on similar topics, appears in the following collection(s): Clinical neurology examination http://n.neurology.org/cgi/collection/clinical_neurology_examination Cluster headache http://n.neurology.org/cgi/collection/cluster_headache EMG http://n.neurology.org/cgi/collection/emg Genetic linkage http://n.neurology.org/cgi/collection/genetic_linkage Neuropathic pain http://n.neurology.org/cgi/collection/neuropathic_pain Peripheral neuropathy http://n.neurology.org/cgi/collection/peripheral_neuropathy |
| Permissions & Licensing | Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions |
| Reprints | Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise |

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2022 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

