hemolytic streptococcal infection. J Child Neurol 1998;13: 354–356.

- Muffat-Joly M, Barry B, Henin D, Fay M, Gehanno P, Pocidalo JJ. Otogenic meningoencephalitis induced by *Streptococcus pneumoniae* in gerbils. Arch Otolaryngol Head Neck Surg 1994;120:925–930.
- Blanot S, Muffat-Joly M, Vilde F, et al. A gerbil model for rhombencephalitis due to Listeria monocytogenes. Microb Pathog 1997;23:39-48.
- Watanabe-Ohnishi R, Aelion J, leGros L, et al. Characterization of unique human TCR V beta specificities for a family of streptococcal superantigens represented by rheumatogenic serotypes of M protein. J Immunol 1994;152:2066-2073.
- 32. Toyosaki T, Yoshioka T, Tsuruta Y, Yutsudo T, Iwasaki M, Suzuki R. Definition of the mitogenic factor (MF) as a novel streptococcal superantigen that is different from streptococcal pyrogenic exotoxins A, B, and C. Eur J Immunol 1996;26: 2693-2701.
- 33. Rikiishi H, Okamoto S, Sugawara S, Tamura K, Liu ZX, Kumagai K. Superantigenecity of helper T cell mitogen (SPM-2) isolated from culture supernatants of streptococcus pyogenes. Immunology 1997;91:406-413.
- Vandevyver C, Mertens N, van den Elsen P, Medaer R, Raus J, Zhang J. Clonal expansion of myelin basic protein-reactive

T cells in patients with multiple sclerosis: restricted T cell receptor V gene rearrangements and CDR3 sequence. Eur J Immunol 1995;25:958–968.

- Maillard I, Luth F, Acha-Orbea H, Diggelman H. Role of the immune response induced by superantigens in the pathogenesis of microbial infections. Parasitol 1997;115:S67–S78.
- Herrmann T, Baschieri S, Less RK, MacDonald HR. In vivo responses of CD4+ and CD8+ cells to bacterial superantigens. Eur J Immunol 1992;22:1935–1938.
- 37. Herrmann T, MacDonald HR. The CD8 T cell response to staphylococcal enterotoxins. Semin Immunol 1993;5:33-39.
- Hermans G, Denzer U, Lohse A, Raus J, Stinissen P. Cellular and humoral immune response against autoreactive T cells in multiple sclerosis patients after T cell vaccination. J Autoimmun 1999;13:233–246.
- Pohl-Koppe A, Burchett SK, Thiele EA, Hafler DA. Myelin basic protein reactive Th2 T cells are found in acute disseminated encephalomyelitis. J Neuroimmunol 1998;91:19-27.
- Genain CP, Abel K, Belmar N, et al. Late complications of immune deviation therapy in a nonhuman primate. Science 1996;274:2054-2057.
- Franchimont D, Louis E, Dewe W, et al. Effects of dexamethasone on the profile of cytokine secretion in human whole blood cell cultures. Regul Pept 1998;73:59-65.

Neuro Images




Figure. MRI at admission (postcontrast T1-weighted). (A) Ring-enhancing left cerebellar lesion and meningeal contrast enhancement. (B) Multiple intramedullary lesions.

Cerebellar and medullar histoplasmosis

M.J. Vos, MD, Y.J. Debets-Ossenkopp, MD, PhD, F.A.P. Claessen, MD, G.J. Hazenberg, MD, J.J. Heimans, MD, PhD, Amsterdam, the Netherlands

A 16-year-old Surinam boy presented with a 3-month history of progressive paraparesis and periods of fever and neck stiffness. Neurologic examination demonstrated a severe spastic paraparesis, Babinski signs, and sensory disturbance under level T4. The patient was not taking immunosuppressants, was HIV-seronegative, and had normal CD4⁺ lymphocyte counts. Postcontrast T1-weighted MRI of brain and spinal cord revealed a ring-enhancing left cerebellar lesion, meningeal contrast enhancement, and multiple intramedullary lesions (figure). CSF examination showed a lymphocytic pleocytosis, elevated total protein, lowered glucose, and negative Gram staining and cultures. Biopsy of one intramedullary lesion disclosed granulomatous lymphohistiocytic inflammation with giant cells and necrosis; cultures were positive for *Histoplasma capsulatum*. Some neurologic improvement occurred after sequential treatment with amphotericin B and itraconazole.

Primary neurologic presentation of histoplasmosis is extremely rare, especially in young, nonimmunocompromised hosts. Tan et al. reported a case of disseminated histoplasmosis in a 72-year-old man, which was the first report of histoplasmosis presenting as a myelopathy.¹ Tabbal and Harik recently reported cerebral histoplasmosis in a 60-yearold man, presenting as a bilateral pyramidal syndrome.²

To our knowledge, this is the first case of cerebellar, meningeal, and medullar histoplasmosis reported in a young, nonimmunocompromised host. The severity of the neurologic presentation, lack of systemic manifestations, and extensiveness of CNS lesions on MRI are exceptional.

Tan V, Wilkins P, Badve S, et al. Histoplasmosis of the central nervous system. JNNP 1992;55:619–622.

Tabbal SD, Harik SI. Cerebral histoplasmosis. N Engl J Med 1999;340: 1176.

Neurology®

Cerebellar and medullar histoplasmosis Neurology 2000;54;1441 DOI 10.1212/WNL.54.7.1441

This information is current as of April 11, 2000

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/54/7/1441.full
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/54/7/1441.full##otherarticles
Permissions & Licensing	Information about reproducing this article in parts (figures,tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology ® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

