

21. Schifter S. Circulating concentrations of calcitonin gene-related peptide (CGRP) in normal man determined with a new, highly sensitive radioimmunoassay. *Peptides* 1991;12:365–369.
22. Buzzi MG, Carter WB, Shimizu T, Heath H, Moskowitz MA. Dihydroergotamine and sumatriptan attenuate levels of CGRP in plasma in rat superior sagittal sinus during electrical stimulation of the trigeminal ganglion. *Neuropharmacology* 1991;30:1193–1200.
23. Levine JD, Fields HL, Basbaum AI. Peptides and the primary afferent nociceptor. *J Neurosci* 1993;13:2273–2286.
24. Morton CR, Hutchison WD. Morphine does not reduce the intraspinal release of calcitonin gene-related peptide in the cat. *Neurosci Lett* 1990;117:319–324.
25. Zaidi M, Bevis PJ, Abeyasekera G, et al. The origin of circulating calcitonin gene-related peptide in the rat. *J Endocrinol* 1986;110:185–190.
26. Wimalawansa SJ, MacIntyre I. Heterogeneity of plasma calcitonin gene-related peptide: partial characterisation of immunoreactive forms. *Peptides* 1988;9:407–410.
27. Goadsby PJ, Edvinsson L, Ekman R. Release of vasoactive peptides in the extracerebral circulation of humans and the cat during activation of the trigeminovascular system. *Ann Neurol* 1988;23:193–196.
28. Goadsby PJ, Edvinsson L. The trigeminovascular system and migraine: studies characterizing cerebrovascular and neuropeptide changes seen in humans and cats. *Ann Neurol* 1993;33:48–56.
29. Gallai V, Sarchielli P, Floridi A, et al. Vasoactive peptide levels in the plasma of young migraine patients with and without aura assessed both interictally and ictally. *Cephalgia* 1995;15:384–390.
30. Ekins RP, Newman GB. Optimization of radioimmunoassay, protein binding and other saturation assay techniques. *J Endocrinol* 1973;58:xxi.
31. Bach FW, Langemark M, Ekman R, Rehfeld JF, Schifter S, Olesen J. Effect of sulphiride or paroxetine on cerebrospinal fluid neuropeptide concentrations in patients with chronic tension-type headache. *Neuropeptides* 1994;27:129–136.
32. Girgis SI, Macdonald DW, Stevenson JC, et al. Calcitonin gene-related peptide: potent vasodilator and major product of calcitonin gene. *Lancet* 1985;6:14–16.

NeuroImages

MR choroid plexus sign of iron overload

R. Kira, MD, PhD, S. Ohga, MD, PhD, H. Takada, MD, PhD, K. Gondo, MD, PhD, F. Mihara MD, PhD, T. Hara, MD, PhD, Fukuoka, Japan

A 7-year-old boy with Diamond–Blackfan anemia was transfusion-dependent at age 4, at which point hemochromatosis progressed. Laboratory studies showed the following: iron 292 $\mu\text{g/dL}$ (normal, 70 to 190 $\mu\text{g/dL}$), unsaturated iron binding capacity 29 $\mu\text{g/dL}$ (normal, 130 to 370 $\mu\text{g/dL}$), and ferritin 6403 ng/mL (normal, 26.0 to 250.0 ng/mL). A brain MR study indicated the restricted accumulation of iron in the choroid plexus (figure). CT demonstrated no calcification on the choroid plexus. These findings are compatible with neuropathologic examination of a patient with hemochromatosis.¹

According to the autoradiographic distribution in the adult rat brain, the choroid plexus showed the highest uptake level of ⁵⁹Fe from the blood.² Thus, the choroid plexus might protect the brain from iron overloading through a buffering effect.

MRI, especially T2*-weighted imaging, effectively detects iron overloading because of a striking reduction of T2 relaxation time. Further investigations by using T2*-weighted imaging are necessary in patients with various causes of iron overload including primary hemochromatosis and acute iron intoxication.




Figure. (A) Axial T2-weighted MRI shows decreased signal intensity over the bilateral choroid plexus (arrows). (B) A corresponding T2-weighted image reveals more decreased and widespread signal intensity (arrows). The choroid plexus appeared normal on T1-weighted imaging and CT scanning.*

References

1. Dwork AJ. Effects of diet and development upon the uptake and distribution of cerebral iron. *J Neurol Sci* 1995;134(suppl):45–51.
2. Morris CM, Keith AB, Edwardson JA, Pullen RG. Uptake and distribution of iron and transferrin in the adult rat brain. *J Neurochem* 1992;59:300–306.

Neurology[®]

MR choroid plexus sign of iron overload

Neurology 2000;55:1340

DOI 10.1212/WNL.55.9.1340

This information is current as of November 14, 2000

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/55/9/1340.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

