


Congestive radiculopathy

Figure T1-weighted sagittal MRI, T2-weighted parasagittal MRI, and coronal reconstructed CT



T1-weighted sagittal MRI: enlarged lumbar epidural veins (arrow) compressing the dural sac (A). T2-weighted parasagittal MRI: enlarged lumbar veins entering the neuroforamina (arrow) in TH12-L5 (B). Coronal reconstructed CT: extensive thrombosis of the inferior vena cava (arrow). Hypodense intrahepatic metastatic lesion (arrowheads) (C).

A previously healthy 34-year-old man was referred for suspected lumbar canal stenosis. He presented with a 2-month history of exercise-induced lower back pain radiating to the thighs and a sensation of constriction in the lower abdomen, readily subsiding in the supine position. Neurologic examination was normal. MRI showed engorged venous vessels but no lumbar stenosis or disc protrusion (figure). He subsequently developed jaundice and generalized pruritus. Abdominal imaging revealed large intrahepatic lesions compressing an extensively thrombosed inferior vena cava. Histologic diagnosis was moderately differentiated adenocarcinoma. Congestive radiculopathy should be considered especially in younger patients with pain localization atypical of lumbar canal stenosis.¹

Simon N. Jacob, MD, Till Diergarten, MD, and Arthur Melms, MD, Tübingen, Germany

Disclosure: The authors report no conflicts of interest.

Address correspondence and reprint requests to Dr. Simon N. Jacob, Department of Neurology, University of Tübingen Medical School, Hoppe-Seyler Str. 3, 72076 Tübingen, Germany; simon.jacob@klinikum.uni-tuebingen.de

1. Paksoy Y, Gormus N. Epidural venous plexus enlargements presenting with radiculopathy and back pain in patients with inferior vena cava obstruction or occlusion. *Spine* 2004;29:2419–2424.

Neurology[®]

Congestive radiculopathy

Simon N. Jacob, Till Diergarten and Arthur Melms

Neurology 2008;70;734

DOI 10.1212/01.wnl.0000302183.86651.e6

This information is current as of February 25, 2008

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/70/9/734.full
References	This article cites 1 articles, 0 of which you can access for free at: http://n.neurology.org/content/70/9/734.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://n.neurology.org/cgi/collection/all_clinical_neurology All Oncology http://n.neurology.org/cgi/collection/all_oncology All Spinal Cord http://n.neurology.org/cgi/collection/all_spinal_cord MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

