

Disclosure: Dr. Simpson has received speaker honoraria and research support from Allergan, Merz, and Solstice, Inc., and performs botulinum toxin injections. Dr. Gracies has received speaker honoraria and research support from Allergan, Merz, and Solstice, Inc. Dr. Graham has received speaker honoraria and research support from Allergan and performs botulinum toxin injections. Dr. Miyasaki has received research support from Boehringer Ingelheim, Huntington Study Group, NIH, Solvay, Solstice, and Teva. Dr. Naumann has received speaker honoraria from Ipsen and Allergan and performs botulinum toxin injections. Dr. Russman has received research support from Allergan and performs botulinum toxin injections. Dr. L. Simpson has received research support from Allergan. Dr. So holds financial interest in Satoris Inc., and has received research support from NIH, Pfizer, Inc., and NeurogesX, Inc.

Copyright © 2009 by AAN Enterprises, Inc.

1. Simpson DM, Gracies J-M, Graham HK, et al. Assessment: Botulinum neurotoxin for the treatment of spasticity (an evidence-based review): Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology* 2008;70:1691–1698.
2. Dietz V, Sinkjaer T. Spastic movement disorder: impaired reflex function and altered muscle mechanics. *Lancet Neurol* 2007;6:725–733.
3. Lance JW. Symposium synopsis. In: Feldman RG, Young RR, Koella WP, ed. *Spasticity: Disordered Motor Control*. Chicago: Year Book; 1980:485.
4. Twitchell TE. The restoration of motor function following hemiplegia in man. *Brain* 1951;74:443–480.
5. Landau WM, Weaver RA, Hornbein TF. Fusimotor nerve function in man: Differential nerve block studies in normal subjects and in spasticity and rigidity. *Arch Neurol* 1961;3:10–23.
6. Landau WM. Clinical neuromyology II: parables of palsy pills and PT pedagogy: a spastic dialectic. *Neurology* 1988;38:1496–1499.
7. Landau WM, Hunt CC. Dorsal rhizotomy, a treatment of unproven efficacy. *J Child Neurol* 1990;5:174–178.
8. Brashear A, Gordon MF, Elovic E, et al. Intramuscular injection of botulinum toxin for the treatment of wrist and finger spasticity after a stroke. *N Engl J Med* 2002;347:395–400.
9. Snow BJ, Tsui JK, Bhatt MH, Varelas M, Hashimoto SA, Calne DB. Treatment of spasticity with botulinum toxin: a double-blind study. *Ann Neurol* 1990;28:512–515.
10. Gracies JM, Lugassy M, Weisz D, et al. Botulinum toxin dilution and endplate targeting in spasticity: a double-blind controlled study. *Arch Phys Med Rehabil* 2009;90:9–16.
11. Gracies JM. Pathophysiology of spastic paresis: part II: the emergence of muscle overactivity. *Muscle Nerve* 2005;31:552–571.

12. Simpson DM, Gracies JM, Yablon SA, Barbano R, Brashear A. Botulinum neurotoxin vs tizanidine in upper limb spasticity: a placebo-controlled study. *J Neurol Neurosurg Psychiatry* Epub 2008 Oct 31.
13. Pak S, Patten C. Strengthening to promote functional recovery poststroke: an evidence-based review. *Top Stroke Rehabil* 2008;15:177–199.

EXCESSIVE ACUTE MIGRAINE MEDICATION USE AND MIGRAINE PROGRESSION

To the Editor: We read with interest the article by Bigal et al.¹ commenting on migraine progression due to excessive use of acute migraine medication.

Transformed migraine is common, invariably disabling, and frequently hard to treat. The authors mention that among acute migraine treatments, opiates and barbiturates are associated with progression of migraine to chronic migraine. The critical dose of exposure for opiates is only 8 days per month and 5 days per month for barbiturates, which demonstrates that these medications are best avoided in the acute treatment of migraine.

Bigal et al. found nonsteroidal anti-inflammatory drugs (NSAIDs) protective in those with <10 days of headache. Our experience with migraine patients has been that they frequently underestimate and understate their frequency and duration of NSAID use. NSAIDs are widely available without prescription. They are the most frequently and widely used medications by patients and doctors for acute migraine treatment. In our experience, transformation of episodic migraine to chronic migraine frequently occurs in the setting of excessive NSAID use.

We would stress that migraine pathophysiology is complex and further investigation is needed. In addition, haphazard use of ibuprofen for headache may cause more harm than good.

Nitin K. Sethi, MD, New York, NY

Disclosure: The author reports no disclosures.

Editor's Note: The authors of the article were offered the opportunity to respond but declined.

Copyright © 2009 by AAN Enterprises, Inc.

1. Bigal ME, Lipton RB. Excessive acute migraine medication use and migraine progression. *Neurology* 2008;71:1821–1828.

CORRECTION

Fatal congenital myopathy and gastrointestinal pseudo-obstruction due to *POLG1* mutations

In the article "Fatal congenital myopathy and gastrointestinal pseudo-obstruction due to *POLG1* mutations" by C. Giordano et al. (*Neurology*® 2009;72:1103–1105), there is an error in the capitalization of an author's name. The author should have been listed as M. De Curtis. The authors regret the error.

Neurology[®]

Fatal congenital myopathy and gastrointestinal pseudo-obstruction due to *POLG1* mutations

Neurology 2009;73;738-738-a
DOI 10.1212/WNL.0b013e3181b7c6e1

This information is current as of August 31, 2009

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/73/9/738.2.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

