

Practice Parameter update: Evaluation and management of driving risk in dementia

Report of the Quality Standards Subcommittee of the American Academy of Neurology



D.J. Iverson, MD
G.S. Gronseth, MD
M.A. Reger, PhD
S. Classen, PhD, MPH,
OTR/L
R.M. Dubinsky, MD,
MPH
M. Rizzo, MD

Address correspondence and
reprint requests to American
Academy of Neurology, 1080
Montreal Avenue, St. Paul, MN
55116
guidelines@aan.com

ABSTRACT

Objective: To review the evidence regarding the usefulness of patient demographic characteristics, driving history, and cognitive testing in predicting driving capability among patients with dementia and to determine the efficacy of driving risk reduction strategies.

Methods: Systematic review of the literature using the American Academy of Neurology's evidence-based methods.

Recommendations: For patients with dementia, consider the following characteristics useful for identifying patients at increased risk for unsafe driving: the Clinical Dementia Rating scale (Level A), a caregiver's rating of a patient's driving ability as marginal or unsafe (Level B), a history of crashes or traffic citations (Level C), reduced driving mileage or self-reported situational avoidance (Level C), Mini-Mental State Examination scores of 24 or less (Level C), and aggressive or impulsive personality characteristics (Level C). Consider the following characteristics not useful for identifying patients at increased risk for unsafe driving: a patient's self-rating of safe driving ability (Level A) and lack of situational avoidance (Level C). There is insufficient evidence to support or refute the benefit of neuropsychological testing, after controlling for the presence and severity of dementia, or interventional strategies for drivers with dementia (Level U). *Neurology*® 2010;74:1316-1324

GLOSSARY

AAN = American Academy of Neurology; **AD** = Alzheimer disease; **CDR** = Clinical Dementia Rating; **CI** = confidence interval; **MMSE** = Mini-Mental State Examination; **OR** = odds ratio; **ORDT** = on-road driving test; **QSS** = Quality Standards Subcommittee; **RR** = relative risk.

Driving skills deteriorate with increasing dementia severity.¹ While patients with mild dementia, as a group, are higher-risk drivers,¹ more recent studies²⁻⁴ report that as many as 76% are still able to pass an on-road driving test (ORDT) and can safely drive.³ Faced with these facts, clinicians caring for patients with dementia seek to identify those patients with cognitive impairment who may be at higher risk for unsafe driving, without unnecessarily restricting those who are safe drivers.

Clinicians' predictions of driving performance, when based primarily on the Mini-Mental State Examination (MMSE), result in no correlation⁵ or have a relatively low sensitivity for identifying an unsafe driver.³ When elements of the driving history and additional cognitive

testing are considered along with MMSE scores, the predictions are more accurate,⁶ but still maintain only moderate degrees of sensitivity and specificity.

This parameter is an update of the 2000 American Academy of Neurology (AAN) practice parameter on driving and dementia. In addition, this parameter seeks to identify historical features that are associated with increased driving risk.

The parameter addresses the following clinical questions:

1. How strongly are global measures of dementia severity associated with decreased driving ability?
2. To what extent are patients and their caregivers able to assess driving ability and risk?

Supplemental data at
www.neurology.org

e-Pub ahead of print on April 12, 2010, at www.neurology.org.

From the Humboldt Neurological Medical Group, Inc. (D.J.I.), Eureka, CA; Department of Neurology (G.S.G., R.M.D.), University of Kansas, Kansas City; National Center for Telehealth and Technology (M.A.R.), Defense Centers of Excellence for Psychological Health and Traumatic Brain Injury, Tacoma, WA; Occupational Therapy Department (S.C.), University of Florida, Gainesville; and Department of Neurology (M.R.), University of Iowa, Iowa City.

Appendices e-1–e-5, tables e-1–e-3, and references e1–e10 are available on the *Neurology*® Web site at www.neurology.org.

Approved by the Quality Standards Subcommittee on November 5, 2008; by the Practice Committee on February 5, 2009; and by the AAN Board of Directors on December 14, 2009.

Disclosure: Author disclosures are provided at the end of the article.

3. Which elements of the driving history are associated with decreased driving ability?
4. Which neuropsychological tests provide additional prognostic information?
5. Are there any interventions that reduce driving risk?

DESCRIPTION OF THE ANALYTIC PROCESS

Panel formation and literature search. The AAN invited neurologists, a neuropsychologist, an occupational therapist with content domain expertise, and neurologists with methodologic expertise to perform this review. Relevant articles published between 1970 and December 2006 were identified using the search strategy listed in appendix e-3 on the *Neurology*[®] Web site at www.neurology.org.

Assessing the evidence. All eligible articles were rated by one Quality Standards Subcommittee (QSS) member and one author panel member, using the 4-tiered scheme for rating a diagnostic study's risk of bias (appendix e-4). Differences between reviewers were adjudicated by a second QSS member. Recommendations were linked to the strength of the evidence (appendix e-5).

The panel reviewed studies of patients with dementia of any cause or mild cognitive impairment. Population studies of aged drivers without an a priori diagnosis of dementia were accepted for analysis when studies limited to drivers with dementia were unavailable or inconclusive. The justification for this is based on the strong correlation between aging and dementia, and the fact that these studies frequently identified individuals with cognitive impairment without a previous diagnosis of dementia.⁷⁻¹⁰ To account for spectrum bias, such studies were downgraded by one evidence class, per QSS precedent.

Accepted outcome measures included ORDT performance, driving simulator performance, crash data, and caregiver reports. The panel considered standardized ORDTs to be the most valid measure. State-administered ORDTs are the de facto legal determinant of driving ability. Driving simulator studies can evaluate driving behavior in (simulated) dangerous circumstances, but have varying degrees of standardization and validity.¹¹ Crash data are somewhat insensitive, because not all certifiably unsafe drivers have had a crash, and nonspecific, because not all drivers with an at-fault crash are unsafe drivers.

Cohort studies were judged to have a narrow spectrum if they excluded subjects based upon the value of a predictor variable (e.g., a study of the predictive ability of the MMSE that excluded subjects with MMSE scores less than 25). Case-control studies were judged to have a narrow spectrum if they excluded equivocal outcomes (e.g., a study of at-fault

crashes that excluded crashes with indeterminate at-fault status).

ANALYSIS OF EVIDENCE Of approximately 6,000 studies identified by the search strategy, 422 were selected for full-text review. A secondary bibliography search yielded 80 additional references.

The evidence tables (tables e-1 through e-3) list the characteristics, class of evidence, and results of some of the studies reviewed. When multiple studies addressing the same question were available, only the studies with lowest risk of bias (Class I or II) are discussed in the text.

How strongly are global measures of dementia severity associated with decreased driving ability? *Clinical Dementia Rating.* In a Class I study,³ relative risk (RR) for failing an ORDT was 82.7 (confidence interval [CI] 5.1–1333) for Clinical Dementia Rating (CDR) 0.5 (table) and 88.67 (CI 5.4–1444) for CDR 1, compared to drivers with CDR 0 (note wide CIs). However, 85% of the CDR 0.5 group and 76% of the CDR 1 group still passed the ORDT, compared to 100% of CDR 0. In a second Class I study,¹² drivers with a CDR of 1 were more likely to be judged unsafe on 6-month follow-up ORDTs than drivers with a CDR of 0 (RR 2.68, CI 1.4–4.8; NS for CDR of 0.5 vs 0 and CDR 1 vs 0.5). A Class II study¹³ reported that drivers with a CDR of 0.5 had an RR of 9.67 (CI 2.3–40.7) for being judged unsafe in comparison to drivers with a CDR of 0; drivers with a CDR of 1 had an RR of 12 (CI 2.8–50.1). Yet 67% of the CDR 0.5 group and 41% of the CDR 1 group still passed the ORDT, vs 78% of controls. In a second Class II study,² drivers with a CDR of 0.5 or 1 had an RR for unsafe driving of 25 (CI 1.5–384) compared to drivers with a CDR of 0; however, there was no difference in CDR between safe and unsafe drivers ($p = 0.90$), and 45% of drivers with a CDR of 0.5–1 passed the ORDT.

Conclusions. The CDR is established as useful for identifying patients at increased risk for unsafe driving (2 Class I and 2 Class II studies); however, a substantial number of patients with a CDR of 0.5–1 (41%–85%) will be found to be safe drivers by an ORDT.

MMSE. Studies evaluating the MMSE have reported conflicting results. In a Class II study,¹⁴ MMSE was strongly correlated with ORDT score ($r = 0.63$). In a second Class II study,⁵ 64% of subjects with an MMSE of 24 or less failed an ORDT, and MMSE was significant in logistic regression modeling. In a Class III study,¹⁵ MMSE was also correlated with ORDT scores ($r = 0.72, p < 0.01$).

However, in a Class II study,² MMSE score (mean = 24) was not correlated with ORDT score.

Table Clinical Dementia Rating^a

Categories	Impairment			
	0	0.5	1	2
Memory (major category)	No memory loss or slight inconsistent forgetfulness	Consistent slight forgetfulness, partial recollection of events, "benign" forgetfulness	Moderate memory loss; more marked for recent events; defect interferes with everyday activities	Severe memory loss; only highly learned material retained; new material rapidly lost
Secondary categories				
Orientation	Fully oriented	Fully oriented except for slight difficulty with time relationships	Moderate difficulty with time relationships; oriented for place at examination; may have geographic disorientation elsewhere	Severe difficulty with time relationships; usually disoriented to time, often to place
Judgment and problem solving	Solves everyday problems and handles business and financial affairs well; judgment good in relation to past performance	Slight impairment in solving problems, similarities, and differences	Moderate difficulties in handling problems, similarities, and differences; social judgment usually maintained	Severely impaired in handling problems, similarities, and differences; social judgment usually impaired
Community affairs	Independent function at usual level in job, shopping, and volunteer and social groups	Slight impairment in these activities	Unable to function independently at these activities although may still be engaged in some; appears normal to casual inspection	No pretense of independent function outside home; appears well enough to be taken to function outside a family home
Home and hobbies	Life at home, hobbies, and intellectual interests are well-maintained	Life at home, hobbies, and intellectual interests slightly impaired	Mild but definite impairment in function at home, more difficult chores abandoned, more complicated hobbies and interests abandoned	Only simple chores preserved; very restricted interests, poorly maintained
Personal care	Fully capable of self-care	Fully capable of self-care	Needs prompting	Requires assistance in dressing, hygiene, keeping of personal effects

^aImpairment is the decline from the subject's usual level of functioning. Clinical Dementia Rating (CDR) = Memory score unless 3 or more of the secondary categories score above or below the Memory score, in which case the CDR = the majority of the secondary categories. For complete instructions, see Morris.⁴¹

Three additional Class III studies reported no correlation with ORDT scores^{16,17} or crashes.¹⁸

Conclusions. An MMSE score of ≤ 24 is possibly useful in identifying patients at increased risk for unsafe driving (1 Class II study). Otherwise, the correlation between MMSE scores and driving performance is unclear, as data are conflicting.

To what extent are patients and their caregivers able to assess driving ability and risk? Patients who rate themselves as poor or fair drivers have usually either begun to restrict their driving or have stopped entirely (Class II).¹⁹ Those who continue to drive with self-imposed restrictions have a fivefold increased risk of crashes (Class III).¹⁸

In a Class I study of patients with mild Alzheimer disease (AD) (CDR 0.5 to 1),³ 94% rated themselves as safe, but only 41% passed an ORDT (specificity for safe 10.7%). In another Class I study of patients with mild AD,²⁰ all patients who failed the ORDT considered themselves to be safe drivers. A third Class I study of patients with mild AD²¹ also reported significant discrepancies between self-rating as safe and ORDT performance.

Caregiver ratings correlate modestly with ORDT performance. A Class I study of patients with mild AD³ reported a sensitivity of 47.8% and a specificity of 81.8% for a caregiver's rating of marginal or unsafe, vs an experienced neurologist's sensitivity of 60.7%

and specificity of 90.9%. A second Class I study of patients with mild AD²¹ reported that, while there was no significant difference between informant ratings and patient performance in most categories of an ORDT, informants overrated performance in nearly every category.

Conclusions. A caregiver's rating of marginal or unsafe is probably useful in identifying unsafe drivers (1 Class I study). A patient's self-rating of safe is established as not useful for determining that the patient is safe to drive (3 Class I studies).

Which elements of the driving history are associated with decreased driving ability? Crashes and traffic citations. Because of the association across all age groups between crashes or citations and unsafe driving, these events are the basis for the demerit point system used by insurance and licensing agencies to quantify driving risk.^{22,23}

Evidence from studies of dementia. A Class II study¹⁴ of drivers with dementia reported a correlation of -0.38 between crashes and violations (combined) and ORDT score ($p < 0.0001$).

Evidence from studies of aged drivers or mixed populations (downgraded 1 evidence class). A Class III study⁷ of drivers over age 65, which included people with cognitive impairment, reported a correlation between self-reported crashes in the previous 12 months and lower ORDT scores ($r = 0.29$, $p < 0.007$). A second

Class III study²⁴ of drivers over age 60 reported a correlation between a crash in the previous 3 years and a crash in the following 3 years ($r = 0.09-0.11$, $p < 0.0001$).

A history of previous crashes portends a higher risk of future crashes than does the presence of mild dementia alone. A Class II study of drivers aged 55 through 87⁸ reported that the RR of a crash was higher for drivers with a crash in the previous 5 years vs no crash (RR 2.0; CI 1.06–3.79, $p = 0.03$), while a comparison of crash risk for those with and without cognitive impairment yielded no significant difference (RR 1.17; CI 0.61–2.27). A Class III study of drivers over age 65⁹ reported that the odds ratio (OR) for an at-fault crash in the final year of the study was higher for drivers with one or more at-fault crashes in the first 4 years of the study (OR 2.1; CI 1.5–3.0) than for drivers with mild cognitive impairment (OR 0.8; CI 0.5–1.4). Another Class III study of drivers over age 65¹⁰ reported that the OR for a future crash was higher for drivers with a crash in the previous 2 years (OR 2.0; CI 1.1–3.7) than it was for drivers with mild cognitive impairment (RR 0.6; CI 0.3–1.2).

A Class II study of drivers age 79 or older²⁵ reported an OR for an injury crash of 2.1 (CI 1.1–4.1) in the following 2 years among drivers with a citation in the previous 2 years.

A large Class III study²⁴ of drivers over age 60 ($n = 426,408$) reported a correlation between citations in the previous 3 years and in the following 3 years ($r = 0.08-0.10$, $p < 0.0001$). Another large Class III study²³ of a random sample of all-aged drivers ($n = 144,710$) reported that, for drivers over age 60, each traffic citation in the previous 3 years resulted in an additional RR of 0.54–0.77 for a crash in the following 3 years. For drivers over age 70, 2 or more citations result in RR of a crash that exceeds that of any other age group.

Conclusions. A history of a crash in the previous 1 to 5 years or a traffic citation the previous 2 to 3 years is possibly useful in identifying patients with decreased driving ability (1 Class II and 5 Class III studies). A history of a crash is possibly more useful in identifying patients at risk for subsequent crashes than the presence of mild dementia alone (3 Class III studies).

Reduced mileage and situational avoidance. Evidence from studies of aged drivers or mixed populations (downgraded one evidence class). Class I and Class II studies^{2,26,27} have reported that driving mileage is significantly reduced in patients with mild dementia, due to external and self-imposed restrictions. In a Class II mixed-population study²⁸ of people over age 55 and patients with mild AD, the mean weekly mileage of the group failing the

ORDT was 64.6 (SD 51) vs 210 (SD 165) for the passing group ($t = -7.22$, $p < 0.001$). Using logistic regression, reduced mileage was the only predictor of failure on an ORDT ($\chi^2 = 12.84$, $p = 0.0003$). In a Class III study of drivers over age 65⁷ that included drivers with cognitive impairment, reduced mileage was moderately correlated with worse scores on the ORDT (standardized $\beta = 0.281$, $t = 3.185$, $p = 0.002$, $r = 0.334$).

A Class II mixed-population study of drivers over age 60 and medically referred drivers (including cognitively impaired drivers)²⁹ found correlations between a self-report of always avoiding driving in the rain or at night and failing an ORDT ($r = 0.33-0.35$, $p \leq 0.01$). In a Class III study,¹⁸ drivers over 65 who reported changing their driving habits because of safety concerns had an RR of 5.3 (95% CI 0.63–44.63; note CI extends below 1) of a crash in the following 2 years. In a Class III study of aged drivers that included those with cognitive impairment,³⁰ drivers referred to licensing agencies were 3 times as likely to report that they always avoided driving in the rain or at night. However, in a Class II²⁹ and a Class III study,³⁰ drivers who failed an ORDT or had an at-fault crash were just as likely to report never avoiding these situations as reporting always avoiding them.

Conclusions. In mixed-population studies of aged drivers and drivers with mild dementia, reduced driving mileage is possibly associated with an increased risk of poor driving performance (1 Class II, 1 Class III study). In aged drivers, self-reported avoidance is possibly useful to identify drivers at increased risk (1 Class II study). The absence of self-reported avoidance is possibly not useful for identifying safe drivers (1 Class II and 1 Class III study).

Aggressive personality characteristics. In patients with mild to moderate dementia (mean MMSE = 21.9), a Class II study³¹ reported that agitation and aggression were predictive of a refusal to discontinue driving (hazard ratio for driving cessation = 0.54, 95% CI 0.32–0.90). In a Class III study of all-aged drivers (downgraded one evidence class),³² deliberate violations of driving laws, across all ages, correlated with future crashes ($\beta = -0.12$, $p < 0.001$).

Conclusions. Aggressive or impulsive personality characteristics are possibly useful to identify patients with increased driving risk (1 Class II and 1 Class III study).

Which neuropsychological tests provide additional prognostic information? Given the association between global measures of cognitive impairment and driving impairment, one would expect driving impairment to be associated with impairment in individual cognitive domains. Numerous studies confirm

this association with tests including Trails A,⁷ Trails B,^{2,27,33-35} Useful Field of View,^{27,33-35} Line Orientation,^{36,37} Block Design,^{33,36} Benton Visual Retention Test–Copy,²⁰ Benton Visual Retention Test–Recall,^{20,33,35} Complex Figure Test–Copy,^{33,35} Facial Recognition,^{33,35} Logical Memory,^{20,37} and Controlled Oral Word Association.^{27,33,35}

The relevant question for the clinician is whether test results from specific cognitive domains provide any additional value in identifying unsafe drivers beyond dementia diagnosis and severity. This requires studies to control for the presence of dementia (i.e., exclude the control group in the determination of effect size)³⁸ as well as the severity of dementia.³⁹ While the studies referenced above may have controlled for the presence or severity of dementia by stratification or logistic regression, none of the studies adequately controlled for both.

Conclusions. Comprehensive neuropsychological assessment is another means of assessing global cognitive impairment that may be complementary to that of a bedside examination and an informant interview. While neuropsychological testing itself may better define dementia severity, there is insufficient evidence to support or refute the benefit of neuropsychological testing in evaluating driving risk in patients with dementia.

Are there interventions that reduce driving risk? A Class II study⁴⁰ reported that in-person license renewal resulted in a minor reduction in fatal crash risk for drivers over age 85 (RR 0.83, CI 0.72–0.96). Otherwise, there was no evidence to support or refute the use of interventional strategies (e.g., licensing restrictions, driver training) to reduce driving risk. A systematic review^{e1} conducted after our study period was also inconclusive.

Conclusions. There is insufficient evidence to support or refute a benefit of interventional strategies for drivers with dementia.

RECOMMENDATIONS For patients with dementia, consider the following characteristics useful for identifying patients at increased risk for unsafe driving:

- The CDR scale (Level A)
- A caregiver’s rating of a patient’s driving ability as marginal or unsafe (Level B)
- A history of traffic citations (Level C)
- A history of crashes (Level C)
- Reduced driving mileage (Level C)
- Self-reported situational avoidance (Level C)
- MMSE scores of ≤ 24 (Level C)
- Aggressive or impulsive personality characteristics (Level C).

For patients with dementia, consider the following characteristics not useful for identifying patients at increased risk for unsafe driving:

- A patient’s self-rating of safe driving ability (Level A)
- Lack of situational avoidance (Level C)

There is insufficient evidence to support or refute the benefit of neuropsychological testing, after controlling for the presence and severity of dementia, or interventional strategies for drivers with dementia (Level U).

PUTTING THE EVIDENCE IN A CLINICAL CONTEXT Clinicians have professional and, in some cases, legal obligations to identify conditions, such as potentially unsafe driving, that may risk their patients’ or the public’s health. Because there is no test result or historical feature that accurately quantifies driving risk, clinicians are only capable of making qualitative estimates of driving risk.

Clinicians may present patients and their caregivers with the data showing that, as a group, patients with mild dementia (CDR of 1) are at a substantially higher risk for unsafe driving and thus should strongly consider discontinuing driving. At the very least, patients and their caregivers should prepare for the eventuality of driving cessation as dementia severity increases.

However, advocates for maintaining driving privileges may cite the wide CIs for relative risk and ORDT pass rates of 41%¹³ to 76%³ as evidence against a categorical recommendation for patients with mild dementia to cease driving. Such advocates do not want truly capable drivers to negatively impact themselves^{e2,e3} or their caregivers^{e4} by premature driving cessation.

In that case, one may look for evidence of increased risk in an individual patient—concerns from the patient or caregiver, restricted driving, crashes, tickets, and aggressive or reckless driving.^{e5,e6} Consideration of these additional issues can result in a more accurate prediction of driving performance. The questionnaires in Appendices 1 and 2 include these issues and may be useful as a starting point for obtaining pertinent information.

A clinician may wish to integrate this information into an algorithm, such as that in the figure, to obtain a qualitative estimate of driving risk. Given the current state of evidence, this algorithm should only be considered supplementary to the clinician’s judgment.

Patients at higher risk may agree to surrender privileges. For those who wish to continue driving, clinicians may consider referral for a professional or governmental driving evaluation, depending on state

dictors of unsafe driving, 3) report results in terms of relative risk based on the presence of a risk factor or cutoff score, rather than in terms of correlation coefficients,^{e10} and 4) report using STARD-D criteria.

Rules for determining when to revoke driving privileges are ultimately made by society through the legislative process. By accurately quantifying risk with well-designed prospective studies using validated outcome measures, researchers can ensure that legislators are provided with the best science with which to inform their decisions.

ACKNOWLEDGMENT

The authors thank Rudi De Raedt, PhD, and Brian Ott, MD, for performing additional analyses of their original studies; Anthony C. Stein, PhD; Esther Wagner of NHTSA; Thomas Getchius and Laura Moses of the AAN; Lindsay Vaux and David A. Anderson of the University of Iowa for assisting with the literature review; and Reggi Porter and Debra David of the Humboldt Neurological Medical Group for their assistance in the word processing of the manuscript.

DISCLOSURE

Dr. Iverson reports no disclosures. Dr. Gronseth serves as an editorial advisory board member of *Neurology Now*; serves on a speakers' bureau for Boehringer Ingelheim; and receives honoraria from Boehringer Ingelheim and the American Academy of Neurology. Dr. Reger has served on the *Journal of Psychological Training* Review Board and the *Medical Science Monitor* International Reviewers Panel; and has received research support from the Telemedicine and Advanced Technology Research Center AMEDD Advanced Medical Technology Initiative. Dr. Classen has received travel expenses and/or honoraria for lectures or educational activities not funded by industry; and has received research support from the NIH (1R21AG031717-01 [PI]), the VA (00066678 [Co-I]), Florida Department of Transportation (Project Director), and Center for Applications of Psychological Type (PI). Dr. Dubinsky has served on the scientific advisory board and speakers' bureau for Allergan, Inc.; has received honoraria from BrioMed; has received research support from Allergan, Inc., Merz Pharmaceuticals GmbH, and the NIH [NHGRI/NINDS 1R01HG02449-01 (site investigator), NIAM/NINDS R01NS052592 (site investigator), NIAM/NINDS R01NS052619-01 (site investigator), NIAM/NINDS R01NS052592-01 (site investigator), NCCAM 2007P000827 (site investigator), NCCAM U01AT000613 (site investigator)]; and his spouse owns stock in Abbott. Dr. Rizzo has received research support from the NIH [NIA AG17177 (PI), NIA AG026027 (PI), NINDS NS044930 (Co-PI)].

DISCLAIMER

This statement is provided as an educational service of the American Academy of Neurology. It is based on an assessment of current scientific and clinical information. It is not intended to include all possible proper methods of care for a particular neurologic problem or all legitimate criteria for choosing to use a specific procedure. Neither is it intended to exclude any reasonable alternative methodologies. The AAN recognizes that specific patient care decisions are the prerogative of the patient and the physician caring for the patient, based on all of the circumstances involved. The clinical context section is made available in order to place the evidence-based guideline(s) into perspective with current practice habits and challenges. No formal practice recommendations should be inferred.

CONFLICT OF INTEREST

The American Academy of Neurology is committed to producing independent, critical and truthful clinical practice guidelines (CPGs). Significant efforts are made to minimize the potential for conflicts of interests to influence the recommendation of this CPG. To the extent possible, the

AAN keeps separate those who have a financial stake in the success or failure of the products appraised in the CPGs and the developers of the guidelines. Conflict of interest forms were obtained from all authors and reviewed by an oversight committee prior to project initiation. AAN limits the participation of authors with substantial conflicts of interest. The AAN forbids commercial participation in, or funding of, guideline projects. Drafts of the guidelines have been reviewed by at least three AAN committees, a network of neurologists, *Neurology*[®] peer reviewers, and representatives from related fields. The AAN Guideline Author Conflict of Interest Policy can be viewed at www.aan.com.

APPENDIX 1: PATIENT QUESTIONNAIRE

This questionnaire addresses historical features with Level A, Level B, or Level C evidence of relevance to driving competency, as well as selected items from the Manchester Driver Behavior Questionnaire. It is only intended to be used in the qualitative determination of driving risk in elderly patients and patients with dementia, and has not been validated for use in the quantitative determination of driving risk.

1. How many times have you been stopped or ticketed for a traffic violation in the last three years? (0, 1, 2, 3, 4 or more)
2. How many accidents have you been in, or caused, within the last three years? (0, 1, 2, 3, 4 or more)
3. In how many accidents were you at fault in the last three years? (0, 1, 2, 3, 4 or more)

Use this scale to answer the following questions: 1 = strongly disagree; 2 = disagree; 3 = no opinion; 4 = agree; 5 = strongly agree.

1. I have concerns about my ability to drive safely.
2. Others have concerns about my ability to drive safely.
3. I have limited the amount of driving that I do.
4. I avoid driving at night.
5. I avoid driving in the rain.
6. I avoid driving in busy traffic.
7. I will drive faster than the speed limit if I think that I won't be caught.
8. I will run a red light if I think that I won't be caught.
9. I will drive after drinking more alcohol than I should.
10. When I get angry with other drivers, I will honk my horn, gesture, or drive up too closely to them.

How many miles a week do you drive?

APPENDIX 2: FAMILY OR CAREGIVER QUESTIONNAIRE

1. How many times has the patient been stopped or ticketed for a traffic violation in the last three years? (0, 1, 2, 3, 4 or more)
2. How many accidents has the patient been in, or caused, within the last three years? (0, 1, 2, 3, 4 or more)
3. In how many accidents was the patient at fault in the last three years? (0, 1, 2, 3, 4 or more)

Use this scale to answer the following questions: 1 = strongly disagree; 2 = disagree; 3 = no opinion; 4 = agree; 5 = strongly agree.

1. I have concerns about the patient's ability to drive safely.
2. Others have concerns about his/her ability to drive safely.
3. The patient has limited the amount of driving that he/she does.
4. He/she avoids driving at night.
5. He/she avoids driving in the rain.
6. He/she avoids driving in busy traffic.
7. The patient will drive faster than the speed limit if the patient thinks he/she won't be caught.
8. The patient will run a red light if the patient thinks that he/she won't be caught.
9. The patient will drive after drinking more alcohol than the patient should.
10. When he/she gets angry with other drivers, the patient will honk the horn, gesture, or drive up too closely to them.

How many miles a week does the patient drive?

Note: The treating physician must comply with the relevant rules implementing HIPAA, such as 45 C.F.R. § 164.502(g), before disclosing a patient's protected health information to family members or caregivers.

Received February 23, 2009. Accepted in final form December 14, 2009.

REFERENCES

1. Dubinsky RM, Stein AC, Lyons K. Practice parameter: risk of driving and Alzheimer's disease (an evidence-based review). *Neurology* 2000;54:2205–2211.
2. Grace J, Amick MM, D'Abreu A, Festa EK, Heindel WC, Ott BR. Neuropsychological deficits associated with driving performance in Parkinson's and Alzheimer's disease. *J Int Neuropsychol Soc* 2005;11:766–775.
3. Brown LB, Ott BR, Papandonatos GD, Sui Y, Ready RE, Morris JC. Prediction of on-road driving performance in patients with early Alzheimer's disease. *J Am Geriatr Soc* 2005;53:94–98.
4. Ott BR, Heindel WC, Papandonatos GD, et al. A longitudinal study of drivers with Alzheimer's disease. *Neurology* 2008;70:1171–1178.
5. Fox GK, Bowden SC, Bashford GM, Smith DS. Alzheimer's disease and driving: prediction and assessment of driving performance. *J Am Geriatr Soc* 1997;45:949–953.
6. Ott BR, Anthony D, Papandonatos GD, et al. Clinical assessment of the driving competence of patients with dementia. *J Am Geriatr Soc* 2005;53:829–833.
7. De Raedt R, Ponjaert-Kristoffersen I. The relationship between cognitive/neuropsychological factors and car driving performance in older adults. *J Am Geriatr Soc* 2000;48:1664–1668.
8. Owsley C, Ball K, McGwin G Jr, et al. Visual processing impairment and risk of motor vehicle crash among older adults. *JAMA* 1998;279:1083–1088.
9. McGwin G Jr, Sims RV, Pulley L, Roseman JM. Relations among chronic medical conditions, medications, and automobile crashes in the elderly: a population-based case-control study. *Am J Epidemiol* 2000;152:424–431.
10. Foley DJ, Wallace RB, Eberhard J. Risk factors for motor vehicle crashes among older drivers in a rural community. *J Am Geriatr Soc* 1995;43:776–781.
11. Lew HL, Poole JH, Lee EH, Jaffe DL, Huang H-C, Brodd E. Predictive validity of driving-simulator assessments following traumatic brain injury: a preliminary study. *Brain Inj* 2005;19:177–188.
12. Duchek JM, Carr DB, Hunt L, et al. Longitudinal driving performance in early-stage dementia of the Alzheimer type. *J Am Geriatr Soc* 2003;51:1342–1347.
13. Hunt LA, Murphy CF, Carr D, et al. Environmental cueing may affect performance on a road test for drivers with dementia of the Alzheimer type. *Alzheimer Dis Assoc Disord* 1997;11(suppl 1):13–16.
14. Fitten LJ, Perryman KM, Wilkinson CJ, et al. Alzheimer and vascular dementias and driving: a prospective road and laboratory study. *JAMA* 1995;273:1360–1365.
15. Odenheimer GL, Beudet M, Jette AM, et al. Performance-based driving evaluation of the elderly driver: safety, reliability, and validity. *J Gerontol Med Sci* 1994;49:M153–M159.
16. De Raedt R, Ponjaert-Kristoffersen I. Short cognitive/neuropsychological test battery for first-tier fitness-to-drive assessment of older adults. *Clin Neuropsychol* 2001;15:329–336.
17. MacGregor JM, Freeman DH, Zhang D. A traffic sign recognition test can discriminate between older drivers who have and have not had a motor vehicle crash. *J Am Geriatr Soc* 2001;49:466–469.
18. Lesikar SE, Gallo JJ, Rebok GW, Keyl PM. Prospective study of brief neuropsychological measures to assess crash risk in older primary care patients. *JABFP* 2002;15:11–19.
19. Hakamies-Blomqvist L, Wahlström B. Why do older drivers give up driving? *Accid Anal Prev* 1998;30:305–312.
20. Hunt L, Morris JC, Edwards D, Wilson BS. Driving performance in persons with mild senile dementia of the Alzheimer type. *J Am Geriatr Soc* 1993;41:747–753.
21. Wild K, Cotrell V. Identifying driving impairment in Alzheimer disease: a comparison of self and observer reports versus driving evaluation. *Alzheimer Dis Assoc Disord* 2003;17:27–34.
22. Janke MK, Masten SV, McKenzie DM, Gebers MG, Kelsey SL. Teen and senior drivers. California Department of Motor Vehicles; 2003.
23. Gebers MA, Peck RC. The identification of high-risk older drivers through age-mediated point systems. *J Safety Res* 1992;23:81–93.
24. Daigneault G, Joly P, Frigon J-Y. Previous convictions or accidents and the risk of subsequent accidents of older drivers. *Accid Anal Prev* 2002;34:257–261.
25. Keall MD, Frith WJ. Association between older driver characteristics, on-road driving test performance, and crash liability. *Traffic Inj Prev* 2004;5:112–116.
26. Adler G, Kuskowski M. Driving cessation in older men with dementia. *Alzheimer Dis Assoc Disord* 2003;17:68–71.
27. Uc EY, Rizzo M, Anderson SW, et al. Unsafe rear-end collision avoidance in Alzheimer's disease. *J Neurol Sci* 2006;251:35–42.
28. Cushman LA. Cognitive capacity and concurrent driving performance in older drivers. *IATSS Res* 1996;20:38–45.
29. Baldock MRJ, Mathias JL, McLean AJ, Berndt A. Self-regulation of driving and its relationship to driving ability among older adults. *Accid Anal Prev* 2006;38:1038–1045.
30. Staplin L, Lococo KH, Gish KW, Decina L. Model driver screening and evaluation program volume II: Maryland pilot older driver study: DOT HS 809 583. US Department of Transportation; 2003.
31. Herrmann N, Rapoport MJ, Sambrook R, Hébert R, McCracken P, Robillard A. Predictors of driving cessation in mild-to-moderate dementia. *CMAJ* 2006;175:591–595.
32. Parker D, Reason JT, Manstead ASR, Stradling SG. Driving errors, driving violations and accident involvement. *Ergonomics* 1995;38:1036–1048.
33. Rizzo M, Reinach S, McGehee D, Dawson J. Simulated car crashes and crash predictors in drivers with Alzheimer disease. *Arch Neurol* 1997;54:545–551.
34. Uc EY, Rizzo M, Anderson SW, Shi Q, Dawson JD. Driver route-following and safety errors in early Alzheimer disease. *Neurology* 2004;63:832–837.
35. Rizzo M, McGehee DV, Dawson JD, Anderson SN. Simulated car crashes in drivers with Alzheimer disease. *Alzheimer Dis Assoc Disord* 2001;15:10–20.
36. Uc EY, Rizzo M, Anderson SW, Shi Q, Dawson JD. Driver landmark and traffic sign identification in early Al-

- heimer's disease. *J Neurol Neurosurg Psychiatry* 2005;76:764–768.
37. Rebok GW, Keyl PM, Bylsma FW, Blaustein MJ, Tune L. The effects of Alzheimer disease on driving-related abilities. *Alzheimer Dis Assoc Disord* 1994;8:228–240.
38. Reger MA, Welsh RK, Watson GS, Cholerton B, Baker LD, Craft S. The relationship between neuropsychological functioning and driving ability in dementia: A meta-analysis. *Neuropsychology* 2004;18:85–93.
39. Dawson JD, Anderson SW, Uc EY, Dastrup E, Rizzo M. Predictors of driving safety in early Alzheimer disease. *Neurology* 2009;72:521–527.
40. Grabowski DC, Campbell CM, Morrisey MA. Elderly licensure laws and motor vehicle fatalities. *JAMA* 2004;291:2840–2846.
41. Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology* 1993;43:2412–2414.

Neurology[®]

Practice Parameter update: Evaluation and management of driving risk in dementia: Report of the Quality Standards Subcommittee of the American Academy of Neurology

D.J. Iverson, G.S. Gronseth, M.A. Reger, et al.

Neurology 2010;74;1316-1324 Published Online before print April 12, 2010

DOI 10.1212/WNL.0b013e3181da3b0f

This information is current as of April 12, 2010

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/74/16/1316.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2010/04/13/WNL.0b013e3181da3b0f.DC1 http://n.neurology.org/content/suppl/2010/11/18/WNL.0b013e3181da3b0f.DC3 http://n.neurology.org/content/suppl/2010/04/18/WNL.0b013e3181da3b0f.DC2
References	This article cites 39 articles, 7 of which you can access for free at: http://n.neurology.org/content/74/16/1316.full#ref-list-1
Citations	This article has been cited by 8 HighWire-hosted articles: http://n.neurology.org/content/74/16/1316.full##otherarticles
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

