RESIDENT & FELLOW SECTION


Section Editor Mitchell S.V. Elkind, MD, MS

Teaching Neuro*Images*: Neurocutaneous melanosis

Shyamsunder B. Sabat, MD

Address correspondence and reprint requests to Dr. Shyamsunder B. Sabat, 951, 18 St S, Suite 596, Birmingham, AL 35205 ssabat@hmc.psu.edu

Figure Unenhanced and contrast-enhanced MRIs



(A-C) Unenhanced T1-weighted MRI showing hyperintense areas consistent with parenchymal melanin in bilateral amygdala, pons, and cerebellar folia (arrows in A) and opacification of CSF spaces around the lower brainstem and cord (arrows in B, C). (D-F) Contrast-enhanced T1-weighted images showing florid enhancement (arrows) obliterating CSF cisterns at the base of brain and around the entire cord consistent with leptomeningeal melanocytic proliferation.

A 4-year-old boy presented with seizures, dysphagia, and weakness in all extremities. Examination revealed large nevi on the trunk which contained benign melanocytes on biopsy. A diagnosis of neurocutaneous melanosis was made on the basis of the MRI (figure) and CSF examination, which showed numerous melanocytes.

Neurocutaneous melanosis is a nonfamilial neurocutaneous syndrome characterized by large melanocytic nevi and excessive proliferation of melanocytes cells in the leptomeninges. ^{1,2} CNS melanoma develops in about 50%. Patients present with hydrocephalus, seizure, cranial nerve

palsies, intracranial hemorrhage, and myelopathy.¹ Prognosis is poor. Treatment is shunting for the hydrocephalus and supportive.

REFERENCES

- Kadonaga JN, Frieden IJ. Neurocutaneous melanosis: definition and review of the literature. J Am Acad Dermatol 1991;24:747– 755.
- Rhodes RE, Friedman HS, Hatten HP, Hockenberger B, Oakes WJ, Tomita T. Contrast-enhanced MR imaging of neurocutaneous melanosis. AJNR Am J Neuroradiol 1991;12:380–382.

From the University of Alabama at Birmingham, Birmingham. *Disclosure:* The author reports no disclosures.



Teaching NeuroImages: Neurocutaneous melanosis

Shyamsunder B. Sabat *Neurology* 2010;74;e82 DOI 10.1212/WNL.0b013e3181dd4139

This information is current as of May 10, 2010

Updated Information & including high resolution figures, can be found at: **Services** http://n.neurology.org/content/74/19/e82.full

References This article cites 2 articles, 1 of which you can access for free at:

http://n.neurology.org/content/74/19/e82.full#ref-list-1

Subspecialty Collections This article, along with others on similar topics, appears in the

following collection(s): **Developmental disorders**

http://n.neurology.org/cgi/collection/developmental_disorders

MRI

http://n.neurology.org/cgi/collection/mri

Other neurocutaneous disorders

http://n.neurology.org/cgi/collection/other_neurocutaneous_disorders

Permissions & Licensing Information about reproducing this article in parts (figures, tables) or in

its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints Information about ordering reprints can be found online:

http://n.neurology.org/subscribers/advertise

Neurology ® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

