


Teaching Video NeuroImages: Regional myokymia



Yong-Qing Zhang, MD

Address correspondence and reprint requests to Dr. Yong-Qing Zhang, Xiafei Road 123, Xiamen, China yqzhmed@gmail.com

Figure 1 Tibial nerve conduction studies showing consistent F-wave hyperexcitability



A 26-year-old man presented with involuntary muscle twitching of the bilateral lower extremities. The continuous muscle rippling was noted in bilateral calves, posterior thighs, and buttocks (video). Nerve conduction studies revealed F-wave hyperexcitability (figure 1). EMG recording showed doublet, triplet, and multiplet motor unit discharges (figure 2). The clinical presentation and electrophysiologic findings suggested a diagnosis of Isaac syndrome, although


symptoms are typically regional rather than generalized. This patient was successfully treated with phenytoin. Isaac syndrome was first described in 1961,¹ and is characterized by peripheral nerve hyperexcitability which manifests clinically as generalized muscle twitching, stiffness, pseudomyotonia, and cramping. Some patients have muscle hypertrophy and hyperhydrosis. Antibody to voltage-gated potassium channels can be detected in 40% of patients.

Supplemental data at
www.neurology.org

From the Neurology Department, Xiamen Chang Gung Hospital, Xiamen, China.

Disclosure: The author reports no disclosures.

Figure 2 Spontaneous continuous irregularly occurring doublet, triplet, and multiplet single and partial motor unit discharges in medial gastrocnemius



Associated autoimmune disorders and other autoantibodies are found in approximately 50%, most notably antiacetylcholine receptor antibodies, indicating coexistent myasthenia gravis in about 20%. Typical EMG findings include continuous single motor unit discharges occurring as doublets, triplets, and multiplet single unit discharges firing at a high intraburst frequency (150–300 Hz; neuromyotonic discharges), and at lower frequencies (less than 60

Hz; myokymic discharges). The symptoms often respond well to anticonvulsants such as phenytoin and carbamazepine.²

REFERENCES

1. Isaacs H. A syndrome of continuous muscle-fibre activity. *J Neurol Neurosurg Psychiatry* 1961;24:319–325.
2. Maddison P. Neuromyotonia. *J Clin Neurophysiol* 2006; 117:2118–2127.

Neurology[®]

Teaching Video *NeuroImages*: Regional myokymia

Yong-Qing Zhang

Neurology 2010;74:e103-e104

DOI 10.1212/WNL.0b013e3181e2410f

This information is current as of June 7, 2010

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/74/23/e103.full
Supplementary Material	Supplementary material can be found at: http://n.neurology.org/content/suppl/2010/06/06/74.23.e103.DC1
References	This article cites 2 articles, 1 of which you can access for free at: http://n.neurology.org/content/74/23/e103.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Neuromuscular Disease http://n.neurology.org/cgi/collection/all_neuromuscular_disease Autoimmune diseases http://n.neurology.org/cgi/collection/autoimmune_diseases EMG http://n.neurology.org/cgi/collection/emg
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

