RESIDENT & FELLOW SECTION

Section Editor Mitchell S.V. Elkind, MD, MS

Teaching Neuro *Images*: MRI in Ramsay-Hunt syndrome after trigeminal zoster

Donghoon Oh, MD, MS Seong-Ho Park, MD, PhD

Address correspondence and reprint requests to Dr. Seong-Ho Park, Department of Neurology, Seoul National University College of Medicine, Seoul National University Bundang Hospital, 300 Gumi-dong, Bundang-gu, Seongnam-si, Kyeonggi-do, 463-707, South Korea nrpsh@snu.ac.kr

Figure 1 Photograph of patient with right facial mandibular distribution zoster




A 22-year-old man developed painful tongue swelling, a right facial rash, and right lower motor neuron facial palsy over 2 weeks. On examination he had a vesicular rash in a mandibular distribution and on the concha of the right earlobe, and right peripheral facial palsy (figure 1). T2-weighted MRI revealed high signal intensity in the spinal trigeminal tract (figure 2, A and B), and T1-weighted gadoliniumenhanced MRI revealed right facial nerve enhancement in the labyrinthine segment (figure 2C). MRI findings of spinal trigeminal nucleus and tract involvement in trigeminal zoster,1 and of facial nerve enhancement in Ramsay-Hunt syndrome,2 have been reported. Our patient developed trigeminal zoster and Ramsay-Hunt syndrome sequentially, and we confirmed corresponding abnormalities on MRI.

REFERENCES

- Aribandi M, Aribandi L. MRI of trigeminal zoster. Neurology 2005;65:1812.
- Tada Y, Aoyagi M, Tojima H, et al. Gd-DTPA enhanced MRI in Ramsay Hunt syndrome. Acta Otolaryngol Suppl 1994;511:170–174.

Figure 2 MRIs of the patient



Axial T2-weighted MRIs show high signal intensity in the right lower pons and medulla posteriorly and laterally (A, B) (arrows) corresponding to the location of the spinal trigeminal tract. Axial T1-weighted gadolinium-enhanced MRI reveals right facial nerve enhancement in the labyrinthine segment (C) (arrow).

From the Department of Neurology, Seoul National University Bundang Hospital, South Korea. Disclosure: The authors report no disclosures.



Teaching Neuro Images: MRI in Ramsay-Hunt syndrome after trigeminal zoster

Donghoon Oh and Seong-Ho Park Neurology 2010;74;e33 DOI 10.1212/WNL.0b013e3181d25b4a

This information is current as of March 1, 2010

Updated Information & including high resolution figures, can be found at:

Services http://n.neurology.org/content/74/9/e33.full

References This article cites 2 articles, 1 of which you can access for free at:

http://n.neurology.org/content/74/9/e33.full#ref-list-1

Subspecialty Collections This article, along with others on similar topics, appears in the

following collection(s): All Clinical Neurology

http://n.neurology.org/cgi/collection/all_clinical_neurology

MRI

http://n.neurology.org/cgi/collection/mri

Viral infections

http://n.neurology.org/cgi/collection/viral_infections

Permissions & Licensing Information about reproducing this article in parts (figures, tables) or in

its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints Information about ordering reprints can be found online:

http://n.neurology.org/subscribers/advertise

Neurology ® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

