
The psychogenic toe signs

Figure An extended first toe can be due to psychogenic or organic (e.g., striatal toe) dystonia



Passive plantar flexion elicits pain and variable resistance (A). Forced dorsiflexion of the second-fifth toes yields spontaneous plantar flexion of the first toe (B). Conversely, in the case of organic striatal toe, there is no pain or resistance to passive plantar flexion (C) and forced dorsiflexion of the other toes does not alter the spontaneous toe extension (D). (Printed with permission from Mayfield Clinic.)

A 13-year-old boy presented with sudden-onset give-away weakness, nonanatomic sensory loss, and posturing of the right foot followed by episodic left leg tremor, relatively unchanged for 14 months prior to this evaluation. The right first toe was fixed in extension; the others in flexion. Although there was resistance to passive plantar flexion, such movement occurred spontaneously upon forced extension of the second–fifth toes (figure; video on the *Neurology*® Web site at www.neurology.org), unlike the behavior seen in organic striatal toe. Fixed posturing and resistance to manipulation are features of psychogenic dystonia. Foot flexion and inversion, without toe extension, appears to be the more common phenotype of psychogenic foot dystonia.

Alberto J. Espay, MD, MSc, Cincinnati, OH; Anthony E. Lang, MD, FRCPC, Toronto, Canada

Supplemental data at www.neurology.org



Scan this code with your smartphone to access this feature

Author contributions: Dr. Espay: drafting/revising the manuscript, study concept or design, study supervision; Dr. Lang: drafting/revising the manuscript, acquisition of data, study supervision.

Acknowledgment: The authors thank Martha Headworth, Medical illustrator of the UC Neuroscience Institute, for the design and creation of the figure.

Disclosure: Dr. Espay serves/has served on scientific advisory boards for Boehringer Ingelheim, Solvay Pharmaceuticals, Inc., and Abbott; has received honoraria from Novartis, the American Academy of Neurology, and the Movement Disorders Society; serves on the editorial board of *The European Neurological Journal*; serves/has served on the speakers' bureaus for Novartis and UCB; and receives/has received research support from Medtronic, Inc., Allergan, Inc., CleveMed, the University of Cincinnati, the Davis Phinney Foundation, and the Michael J Fox Foundation; and is funded by the KL2 Research Scholars mentored career development award, the NIH Institutional Clinical and Translational Science Award (RR026315-02). Dr. Lang has served on scientific advisory boards for Abbott, Allon Therapeutics, Inc., Biovail Corporation, Boehringer Ingelheim, Cephalon, Inc., Ceregene, Eisai Inc., Medtronic, Inc. Lundbeck Inc., NeuroMolecular Pharmaceuticals, Novartis, Merck Serono, Solvay Pharmaceuticals, Inc., TaroPharma, and Teva Pharmaceutical Industries Ltd.; has received speaker

Copyright © 2011 by AAN Enterprises, Inc.

honoraria from GlaxoSmithKline and UCB; receives/has received research support from the Canadian Institutes of Health Research, the Dystonia Medical Research Foundation, the Michael J. Fox Foundation, the National Parkinson Foundation, and the Ontario Problem Gambling Research Centre; and has served as an expert witness in cases related to the welding industry.

Address correspondence and reprint requests to Dr. Alberto J. Espay, Gardner Center for PD and Movement Disorders, University of Cincinnati Academic Health Center, 260 Stetson St., Suite 2300, Cincinnati, OH 45267-0525; alberto.espay@uc.edu

- Ashour R, Tintner R, Jankovic J. Striatal deformities of the hand and foot in Parkinson's disease. Lancet Neurol 2005;4:423

 431.
- Schrag A, Trimble M, Quinn N, Bhatia K. The syndrome of fixed dystonia: an evaluation of 103 patients. Brain 2004;127: 2360–2372.

NeuroImages Are Free at www.neurology.org!

All Neurology® NeuroImages can now be freely accessed on the Neurology Web site. See them at www.neurology.org, where you can also sign up for journal email alerts and check out other online features, including the Resident & Fellow section, Neurology: Clinical Practice, and the weekly Neurology Podcasts.



The psychogenic toe signs

Alberto J. Espay and Anthony E. Lang *Neurology* 2011;77;508-509 DOI 10.1212/WNL.0b013e318227e4c0

This information is current as of August 1, 2011

Updated Information & including high resolution figures, can be found at:

Services http://n.neurology.org/content/77/5/508.full

Supplementary Material Supplementary material can be found at:

http://n.neurology.org/content/suppl/2011/07/31/77.5.508.DC1

References This article cites 2 articles, 0 of which you can access for free at:

http://n.neurology.org/content/77/5/508.full#ref-list-1

Subspecialty Collections This article, along with others on similar topics, appears in the

following collection(s):

Clinical neurology examination

http://n.neurology.org/cgi/collection/clinical_neurology_examination

Conversion

http://n.neurology.org/cgi/collection/conversion

Dystonia

http://n.neurology.org/cgi/collection/dystonia

Permissions & Licensing Information about reproducing this article in parts (figures, tables) or in

its entirety can be found online at:

 $http://www.neurology.org/about/about_the_journal\#permissions$

Reprints Information about ordering reprints can be found online:

http://n.neurology.org/subscribers/advertise

Neurology ® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2011 by AAN Enterprises, Inc.. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

