



Abstracts

Articles appearing in the April 2018 issue

Neurologist-patient communication about epilepsy in the United States, Spain, and Germany

Background Effective communication between patients and their health care providers is recognized as critically important to improve the quality of health services for individuals with epilepsy. We aimed to describe in-office neurologist-patient conversations about epilepsy and focus on disease identification, shared decision-making, and care planning.



Methods Transcripts and audio recordings of conversations between patients and neurologists in the United States, Spain, and Germany were analyzed linguistically in the topic areas of epilepsy identification and diagnosis, disease education, treatments, and care planning. Analyses included word-level assessments, topic switching, strategies of information elicitation, identification of topics discussed, quantification of questions asked, and assessment of types of questions asked.

Results Conversations of 17 neurologists in the United States, 12 in Spain, and 6 in Germany, with 50, 20, and 16 patients, respectively, were analyzed. Neurologists tended to utilize an event-based, patient-friendly vocabulary to refer to seizures, and in the United States, they avoided using the term "epilepsy." Regardless of who initiated the treatment discussion, the neurologists in all 3 countries were unilaterally responsible for the treatment decision and choice of medication. When describing a new medication, neurologists most often discussed potential side effects but did not review potential benefits. Neurologists rarely defined seizure control and did not ask patients what seizure control meant to them.

Conclusions We identified opportunities related to vocabulary, decision-making, and treatment goal setting that could be targeted to improve neurologist-patient communication about epilepsy, and ultimately, the overall treatment experience and outcomes for patients.

NPub.org/NCP/9101a

SSRI-associated bruxism: A systematic review of published case reports

Purpose of review Antidepressant-associated movement disorders are a well-described phenomenon. However, antidepressant-associated bruxism, jaw pain, or jaw spasm, while reported in dental literature, is less commonly recognized among neurologists. We summarize the clinical features and



treatment of antidepressant-associated bruxism and associated jaw pain through a systematic review of case reports.

Recent findings Antidepressant-associated bruxism may occur in pediatric and adult patients, most commonly among female patients. Patients may develop symptoms with short-term and long-term antidepressant use. Fluoxetine, sertraline, and venlafaxine were the most commonly reported offending agents. Symptoms may begin within 3–4 weeks of medication initiation and may resolve within 3–4 weeks of drug discontinuation, addition of buspirone, or substitution with another pharmacologic agent. The incidence of this phenomenon is unknown.

Summary Bruxism associated with antidepressant use is an underrecognized phenomenon among neurologists, and may be treated with the addition of buspirone, dose modification, or medication discontinuation.

NPub.org/NCP/9101b

Practice Current

Our survey on "When do you suspect autoimmune encephalitis and what is the role of antibody testing?" has received over 1,000 responses from over 80 countries. Explore this topic and others on our redesigned website: compare your practice with peers and see survey results displayed on an interactive world map.

NPub.org/NCP/pc6



What's happening in Neurology® Clinical Practice Neurology 2018;91;33 DOI 10.1212/WNL.00000000005750

This information is current as of July 2, 2018

Updated Information & including high resolution figures, can be found at:

Services http://n.neurology.org/content/91/1/33.full

Permissions & Licensing Information about reproducing this article in parts (figures, tables) or in

its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints Information about ordering reprints can be found online:

http://n.neurology.org/subscribers/advertise

Neurology ® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2018 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

