

Teaching NeuroImages: Substantia nigra T2 hyperintensities in a man with Leber hereditary optic neuropathy


Karl Echiverri, MD, Kyle Darpel, MD, Charles Smith, MD, and Padmaja Sudhakar, MD

Neurology® 2019;93:e1830-e1831. doi:10.1212/WNL.0000000000008458

Correspondence

Dr. Echiverri
Karl.echiverri@uky.edu

Figure MRI scans show symmetric substantia nigra T2 hyperintensities



Coronal T2 MRI (A) and axial fluid-attenuated inversion recovery MRI (B) show bilateral symmetric substantia nigra T2 hyperintensities (arrowheads). These lesions do not show restricted diffusion (arrowheads) on axial diffusion-weighted imaging (C).

A 29-year-old man with alcoholism presented with sudden progressive central visual loss in the right eye. Visual acuity was 20/300 (right) and 20/20 (left). Other findings included right relative afferent pupillary defect and bilateral hyperemic nonedematous optic discs concerning for OD retrobulbar optic neuritis. MRI head and orbit revealed symmetric T2 hyperintensities in the substantia nigra, without optic nerve hyperintensities (figure). Subsequently he lost vision in his left eye. Given that his maternal uncle developed bilateral optic neuritis in his 20s, genetic testing for Leber hereditary optic neuropathy was performed, finding a point mutation (m.11778G>A) in mitochondrially encoded NADH dehydrogenase 4. The unusual gray matter hyperintensities in this case favored a metabolic rather than demyelinating etiology.

MORE ONLINE

→ Teaching slides

links.lww.com/WNL/A990

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

From the Departments of Neurology (K.E., K.D., C.S., P.S.) and Ophthalmology (P.S.), University of Kentucky, Lexington.

Go to Neurology.org/N for full disclosures.

Appendix Authors

Name	Location	Role	Contribution
Karl Echeverri, MD	University of Kentucky	Author	Drafting and revision, literature review
Kyle Darpel, MD	University of Kentucky	Author	Drafting and revision, literature review
Charles Smith, MD	University of Kentucky	Author	Drafting and revision, literature review, imaging acquisition and formatting
Padmaja Sudhakar, MD	University of Kentucky	Author	Drafting and revision

Neurology®

Teaching NeuroImages: Substantia nigra T2 hyperintensities in a man with Leber hereditary optic neuropathy

Karl Echiverri, Kyle Darpel, Charles Smith, et al.

Neurology 2019;93:e1830-e1831

DOI 10.1212/WNL.00000000000008458

This information is current as of November 4, 2019

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/93/19/e1830.full
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Alcohol http://n.neurology.org/cgi/collection/alcohol Mitochondrial disorders http://n.neurology.org/cgi/collection/mitochondrial_disorders MRI http://n.neurology.org/cgi/collection/mri Optic nerve http://n.neurology.org/cgi/collection/optic_nerve Visual loss http://n.neurology.org/cgi/collection/visual_loss
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2019 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

