

Neuropathologic Features of Antemortem Atrophy-Based Subtypes of Alzheimer Disease

Rosaleena Mohanty, PhD, Daniel Ferreira, PhD, Simon Frerich, MSc, J-Sebastian Muehlboeck, MSc, Michel J. Grothe, PhD, and Eric Westman, PhD, on behalf of the Alzheimer's Disease Neuroimaging Initiative

Neurology® 2022;99:e323-e333. doi:10.1212/WNL.0000000000200573

Correspondence

Dr. Mohanty
rosaleena.mohanty@ki.se

Abstract

Background and Objectives

To investigate whether antemortem MRI-based atrophy subtypes of Alzheimer disease (AD) differ in neuropathologic features and comorbid non-AD pathologies at postmortem.

Methods

From the Alzheimer's Disease Neuroimaging Initiative cohort, we included individuals with antemortem MRI evaluating brain atrophy within 2 years before death, antemortem diagnosis of AD dementia/mild cognitive impairment, and postmortem-confirmed AD neuropathologic change. Antemortem atrophy subtypes were modeled as continuous phenomena based on a recent conceptual framework: typicality (spanning limbic-predominant AD to hippocampal-sparing AD) and severity (spanning typical AD to minimal atrophy AD). Postmortem neuropathologic evaluation included AD hallmarks, β -amyloid, and tau as well as non-AD pathologies, alpha-synuclein and TAR DNA-binding protein 43 (TDP-43). We also investigated the overall concomitance across these pathologies. Partial correlations assessed the associations between antemortem atrophy subtypes and postmortem neuropathologic outcomes.

Results

In 31 individuals (26 AD dementia/5 mild cognitive impairment, mean age = 80 years, 26% females), antemortem typicality was significantly negatively associated with neuropathologic features, including β -amyloid ($\rho = -0.39$ overall), tau ($\rho = -0.38$ regionally), alpha-synuclein ($\rho = -0.39$ regionally), TDP-43 ($\rho = -0.49$ overall), and concomitance of pathologies ($\rho = -0.59$ regionally). Limbic-predominant AD was associated with higher Thal phase, neuritic plaque density, and presence of TDP-43 compared with hippocampal-sparing AD. Regionally, limbic-predominant AD showed a higher presence of tau and alpha-synuclein pathologies in medial temporal structures, a higher presence of TDP-43, and concomitance of pathologies subcortically/cortically compared with hippocampal-sparing AD. Antemortem severity was significantly negatively associated with concomitance of pathologies ($\rho = -0.43$ regionally), such that typical AD showed higher concomitance of pathologies than minimal atrophy AD.

MORE ONLINE

 CME Course
NPub.org/cmelist

From the Division of Clinical Geriatrics (R.M., D.F., S.F., J.S.M., E.W.), Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Stockholm, Sweden; Department of Radiology (D.F.), Mayo Clinic, Rochester, MN; Institute for Stroke and Dementia Research (E.W.), University Hospital, Ludwig-Maximilian-University (LMU) Munich, Germany; Unidad de Trastornos del Movimiento (M.J.G.), Servicio de Neurología y Neurofisiología Clínica, Instituto de Biomedicina de Sevilla, Hospital Universitario Virgen del Rocío/CSIC/Universidad de Sevilla, Spain; Clinical Dementia Research Section (M.J.G.), German Center for Neurodegenerative Diseases (DZNE), Rostock, Germany; and Department of Neuroimaging (E.W.), Centre for Neuroimaging Sciences, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Data used in this study were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.ucla.edu). As such, the investigators within the ADNI contributed to the design and implementation of the ADNI and/or provided data but did not participate in the analysis or writing of this report. A complete listing of ADNI investigators can be found in Appendix 2 at links.lww.com/WNL/C7.

The Article Processing Charge was funded by various Swedish foundations including Swedish Foundation for Strategic Research (SSF); the Strategic Research Programme in Neuroscience at Karolinska Institutet (StratNeuro); the Swedish Research Council (VR); the regional agreement on medical training and clinical research (ALF) between Stockholm County Council and Karolinska Institutet; Center for Innovative Medicine (CIMED); the Swedish Alzheimer Foundation; the Swedish Brain Foundation; the Åke Wiberg Foundation; Demensfonden; Stiftelsen Olle Engkvist Byggmästare; Birgitta och Sten Westerberg; Foundation for Geriatric Diseases at Karolinska Institutet; Loo och Hans Ostermans stiftelse för medicinsk forskning; Stiftelsen För Gamla Tjänarinnor; Gun & Bertil Stohnes Stiftelse.

This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Glossary

AD = Alzheimer disease; ADNC = AD neuropathologic change; ADNI = Alzheimer's Disease Neuroimaging Initiative; ADNI-EF = composite cognitive score for executive function; ADNI-MEM = composite scores for memory; aMCI = amnesic mild cognitive impairment; A β = β -amyloid; CDR = Clinical Dementia Rating; MMSC = Mini-Mental State Examination; NCI = neuronal cytoplasmic inclusion; NFT = neurofibrillary tangle; TDP-43 = TAR DNA-binding protein 43; α -syn = α -synuclein.

Discussion

We provide a direct antemortem-to-postmortem validation, highlighting the importance of understanding atrophy-based heterogeneity in AD relative to AD and non-AD pathologies. We suggest that (1) typicality and severity in atrophy reflect differential aspects of susceptibility of the brain to AD and non-AD pathologies; and (2) limbic-predominant AD and typical AD subtypes share similar biological pathways, making them more vulnerable to AD and non-AD pathologies compared with hippocampal-sparing AD, which may follow a different biological pathway. Our findings provide a deeper understanding of associations of atrophy subtypes in AD with different pathologies, enhancing the prevailing knowledge of biological heterogeneity in AD and could contribute toward tracking disease progression and designing clinical trials in the future.

Alzheimer disease (AD) is pathologically defined by the hallmarks of β -amyloid (A β) plaques and tau neurofibrillary tangles (NFTs). However, pure AD is increasingly recognized as not being the most prevalent form of the disease.¹⁻³ Concomitant forms of pathologic proteins such as α -synuclein (α -syn) and TAR DNA-binding protein 43 (TDP-43) have been reported in over 40%⁴ and 50%⁵ of the AD cases, respectively.

Does this multimorbid view of the brain in AD suggest that atrophy may be downstream to not only the AD hallmark pathologies but also the interactions with 1 or more concomitant pathologies? Examination of medial temporal atrophy measured on antemortem MRI in relation to postmortem neuropathology has shown that tau pathology was associated with posterior hippocampal atrophy, whereas TDP-43 pathology was associated with anterior medial temporal atrophy.⁶ Medial temporal atrophy, although a common characteristic, is not always observed in AD. Converging evidence suggests that biological heterogeneity in AD may manifest as distinct atrophy subtypes: typical AD, limbic-predominant AD, hippocampal-sparing AD, and minimal atrophy AD,⁷ with the last 2 showing relatively preserved medial temporal gray matter structure. Thus, revising the initial question, we ask: does this multimorbid view of the brain in AD suggest that atrophy subtypes may be downstream to not only the AD hallmark pathologies but also the interactions with 1 or more concomitant pathologies? To our knowledge, the answer to this question is yet to be explored.

We currently lack in vivo biomarkers to assess pathologies such as α -syn and TDP-43. Therefore, we investigated the relationship between antemortem MRI-based atrophy subtypes and postmortem neuropathologic profiles in AD. Our key research questions are (1) whether antemortem atrophy subtypes of AD are related to individual and/or concomitance of AD and non-AD pathologies at postmortem, and (2) whether this subtype-to-pathology relationship varies by brain

region. Corresponding to these research questions, we hypothesized that antemortem atrophy subtypes of AD may be differentially associated with different AD and non-AD pathologies assessed postmortem, which may vary by brain region.

Methods

Participants

Participants were selected from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (PI: M. Weiner, launched 2003; adni.loni.usc.edu/). The goal of the ADNI is to test and use biomarkers, clinical, and neuropsychological assessments to track disease progression in AD. We included data from participants who had antemortem MRI and postmortem neuropathologic assessments (Version 11, April 12, 2018). eFigure 1, links.lww.com/WNL/C4, shows the selection criteria for this study. Our final cohort comprised 31 participants with intermediate or high AD neuropathologic change (ADNC) at postmortem examination (i.e., pathology-confirmed AD dementia; low ADNC is not an adequate explanation for cognitive impairment or dementia)⁸ and availability of an antemortem MRI scan within 2 years before death (for a more accurate antemortem approximation of the postmortem/final subtype of an individual and to avoid long antemortem-to-postmortem interval being a potential confound).

Standard Protocol Approvals, Registrations, and Patient Consents

All the ADNI protocols were approved by the institutional review boards of each participating institution. All participants provided written informed consent in accordance with the Declaration of Helsinki.

Antemortem Neuroimaging and Cognition

MRI scans were acquired on 1.5T or 3T scanners with T1-weighted sagittal 3D magnetization-prepared rapid gradient-echo

sequences (detailed ADNI imaging protocols: adni.loni.usc.edu/methods/). MRI scans were processed cross-sectionally using FreeSurfer 6.0.0 (freesurfer.net/), automated through the HiveDB system.⁹ Resulting segmentations were visually screened for quality control. Screened scans were included for subsequent analyses. Automatic region of interest parcellation yielded volumes of 41 cortical and subcortical areas^{10,11} per hemisphere, serving as a measure of brain atrophy. We used Mini-Mental State Examination (MMSE),¹² Clinical Dementia Rating (CDR), and composite scores for memory (ADNI-MEM)¹³ and executive function (ADNI-EF)¹⁴ corresponding to the MRI visit as the main outcomes to evaluate the level of cognitive impairment.

Antemortem Atrophy Subtypes

Following the recently proposed conceptual framework for AD subtypes,⁷ we quantified MRI-based atrophy subtypes in terms of 2 principal dimensions: typicality and severity. Given the limited sample size, we modeled atrophy subtypes on a continuous scale for greater sensitivity¹⁵ rather than categorizing individuals into subgroups or categorical subtypes. Typicality was proxied by the ratio of hippocampal volume to whole cortical volume (ratio henceforth referred to as H:C), similar to the index adopted by the original neuropathologic subtyping study.¹⁶ Severity was proxied by the Global Brain Atrophy Index, measured by the ratio of whole brain volume to volume of CSF¹⁷ (ratio henceforth referred to as BV:CSF), such that lower values of the index correspond to more atrophy (i.e., higher severity).

Postmortem Neuropathologic Assessment

Neuropathologic assessments were conducted as part of the ADNI neuropathology core (neuropathologist: Dr. Nigel Cairns, the Knight Alzheimer's Disease Research Center, Washington University School of Medicine, St. Louis, adni.loni.usc.edu/about/#core-container).¹⁸ Assessments followed the NIA-AA guidelines for the neuropathologic assessment of AD⁸ (alz.washington.edu/NONMEMBER/NP/nguide10.pdf).

Antemortem-to-Postmortem Validation Approach

We modeled MRI-based antemortem atrophy subtypes in AD as continuous phenomena¹⁵ of 2 orthogonal typicality and severity dimensions, following the recent conceptual framework for AD subtypes.⁷ We then examined the relationship of these dimensions to postmortem neuropathologic features, including AD (A β , tau) and non-AD (α -syn, TDP-43) pathologies and concomitance across them.

To investigate our first research question of whether antemortem atrophy subtypes of AD may be related to neuropathologic differences, we examined (1) established semiquantitative rating scales for AD-specific neuropathologic measures, including the Thal phase of regional distribution of A β (diffuse and cored) plaques (A0–A3), the Braak stage of NFT distribution (B0–B3), and the Consortium to Establish a Registry for AD scores for the density of neuritic

plaques (C0–C3)⁸; and (2) the presence/absence of comorbid non-AD pathologies, including overall α -syn (Lewy body [LB]) pathology, assessed across the brainstem, limbic region, neocortex, amygdala, and olfactory bulb as per the modified McKeith criteria,^{8,19} and overall TDP-43 pathology assessed as immunoreactive inclusions (comprising any of neuronal cytoplasmic inclusion [NCI], neuronal intraneuronal inclusion, dystrophic neurite, or glial cytoplasmic inclusion) across the amygdala, hippocampus, entorhinal cortex/inferior temporal gyrus, and frontal neocortex.²⁰

To investigate our second research question of whether antemortem atrophy subtypes of AD may be related to postmortem pathologies varying by brain regions, we examined regional pathologic outcomes: we analyzed regions most relevant to atrophy subtypes in AD,⁷ that is, structures of the medial temporal lobe, including the hippocampus at the level of the lateral geniculate nucleus (cornu Ammonis 1 or CA1, dentate gyrus, and parahippocampal gyrus), amygdala, and entorhinal cortex, and structures of the association cortex, including the middle frontal gyrus, superior and middle temporal gyri, and inferior parietal lobe (angular gyrus). We focused on specific forms of pathologies binarized for presence/absence: (1) AD-specific neuropathologic measures of A β (positive for both diffuse and cored plaques) and tau (NFT) and (2) non-AD-specific neuropathologic measures of α -syn (LB) and TDP-43 (NCI).

To investigate whether antemortem atrophy subtypes of AD may be related to concomitance of pathologies that may also vary regionally, we evaluated the total number of pathologies present per region as an outcome: each pathology was binarized for presence/absence and summed, considering both AD-specific and non-AD-specific pathologies (concomitance ranging from 0 through 4).

Statistical Analysis

We analyzed the association between antemortem atrophy subtypes (typicality and severity as continuous independent variables in separate models) and cognition as well as neuropathologic outcomes as dependent variables using linear partial correlations, controlled for age at MRI scan and MRI scanner field strength. Furthermore, each model with typicality as an independent variable was controlled for severity and vice versa to examine whether the correlation may be solely explainable by the dimension treated as independent variable. Because of the limited sample size in this rare antemortem-postmortem dataset, we report significant results at an uncorrected *p* value of <0.05, akin to previous radiologic-pathologic association studies.^{21,22} In addition, we assessed the role of sex (binarized as female or male) and *APOE* status (categorized by all combinations of pairs of the alleles, i.e., 2-4, 3-3, 3-4, and 4-4) through mediation analyses.²³

All statistical analyses and visualizations were conducted using MATLAB R2020b (The MathWorks, Inc., Natick, MA).

Data Availability

Data used in this study have been made publicly available by the ADNI in the Laboratory of Neuro Imaging database.

Results

Participants

Table 1 shows the demographic and antemortem/postmortem characteristics of the cohort. The age at antemortem MRI was 80.0 ± 6.7 years, whereas the age at death was 81.2 ± 6.78 years. The level of cognitive impairment was higher in individuals with AD dementia than those with

Table 1 Characteristics of the Selected Cohort

N	31
Age at antemortem MRI (y)	80.032 ± 6.745
MRI field strength (% 3 tesla)	41.940
Age at death (y)	81.226 ± 6.781
Antemortem MRI to postmortem interval (y)	1.193 ± 0.601
Diagnosis at antemortem MRI	26 AD dementia and 5 aMCI
Sex (% female)	25.810
Education (y)	16.129 ± 2.247
APOE ϵ4 (% carriers)	80.650
MMSE at antemortem MRI	
Overall	18.161 ± 6.738
AD dementia	16.461 ± 5.846
aMCI	27 ± 3.240
CDR at antemortem MRI	
Overall	1.339 ± 0.723
AD dementia	1.500 ± 0.678
aMCI	0.500 ± 0
ADNI-MEM at antemortem MRI	
Overall	-1.268 ± 1.001
AD dementia	-1.515 ± 0.814
aMCI	0.012 ± 0.968
ADNI-EF at antemortem MRI	
Overall	-1.455 ± 1.312
AD dementia	-1.779 ± 1.147
aMCI	0.165 ± 0.802
Presence of markers of cerebrovascular disease postmortem (%)	100

Abbreviations: AD = Alzheimer disease; ADNI-EF = composite cognitive score for executive function; ADNI-MEM = composite cognitive score for memory; aMCI = amnesic mild cognitive impairment; CDR = Clinical Dementia Rating; MMSE = Mini-Mental State Examination.

amnesic mild cognitive impairment (aMCI) in the cohort based on the MMSE, CDR, ADNI-MEM, and ADNI-EF. All individuals had markers of cerebrovascular disease postmortem (1 or more types of the following: macroscopic vascular brain injury, microinfarcts, microbleeds, microhemorrhages, arteriolosclerosis, white matter rarefaction, or other vascular changes).

Antemortem Atrophy Subtypes

Figure 1A shows the atrophy subtypes in antemortem MRI, characterized by the continuous scale measures of typicality (H:C) and severity (BV:CSF). We show 4 examples to illustrate the extremes on each dimension. On the typicality dimension, case RID 1203 represents hippocampal-sparing AD toward the higher extreme, whereas case RID 1393 represents limbic-predominant AD toward the lower extreme. Similarly, on the severity scale, case RID 1271 represents typical AD toward the lower extreme (higher severity), whereas case RID 1425 represents minimal atrophy AD toward the higher extreme (lower severity). The association between typicality and severity was not statistically significant ($r = 0.3$, $p = 0.09$). Antemortem severity ($r = 0.5$, $p = 0.01$; controlled for typicality) but not typicality ($r = -0.1$, $p = 0.6$; controlled for severity) was significantly associated with the MMSE.

Association Between Antemortem Typicality and Neuropathologic Outcomes

Table 2 shows the association between typicality and established neuropathologic rating scales of AD and non-AD pathologies. Most individuals showed a high ADNC at postmortem (Figure 1B). Typicality was significantly associated with Thal A β phase (96.8% at A3, i.e., phase 4–5; Figure 2A), neuritic plaques (87.1% at C3, i.e., frequent neuritic plaques; Figure 2C), and presence of TDP-43 inclusions (Figure 3B). These significant associations were negative, that is, a lower value of H:C (limbic-predominant AD) was associated with a higher pathologic burden or presence of pathology.


Figure 4A shows the association between typicality and regional neuropathologic measures. Typicality was significantly associated with the presence of (1) tau in the dentate gyrus; (2) α -syn in the parahippocampal gyrus; (3) TDP-43 in the parahippocampal gyrus, dentate gyrus, entorhinal cortex, amygdala, and superior/middle temporal gyri; and (4) concomitance of the AD and non-AD pathologies. These associations were negative, that is, a lower value of H:C (limbic-predominant AD) was associated with presence of pathology or higher concomitance of pathologies (eFigure 2–3, [links.ww.com/WNL/C4](https://www.ww.com/WNL/C4)).

Association Between Antemortem Severity and Neuropathologic Outcomes

There were no significant associations between severity and established neuropathologic rating scales of AD and non-AD pathologies (Table 2).

Neither were there any significant associations between severity and regional neuropathologic measures (Figure 4B). However,

Figure 1 Distribution of (A) Antemortem MRI-Based Heterogeneity and (B-C) Postmortem Neuropathology Superposed on MRI-Based Heterogeneity



(A) Antemortem atrophy subtypes modeled as continuous phenomena by the dimensions of typicality and severity. Four individual cases are highlighted, showing the extremes on each dimension; (B) postmortem AD neuropathologic change; and (C) postmortem secondary diagnosis assigned per individual. All plots show antemortem MRI-based typicality on the horizontal scale, proxied by the index = $\left(\frac{\text{hippocampal volume}}{\text{cortical volume}}\right)$; all plots show antemortem MRI-based severity on the vertical scale, proxied by the Global Brain Atrophy Index = $\left(\frac{\text{total brain volume}}{\text{cerebrospinal fluid volume}}\right)$, whereby higher values correspond to lower severity. AD = Alzheimer disease; ADNC = AD neuropathologic change; ALB = amygdala Lewy bodies; aMCI = amnesic mild cognitive impairment; DLB = dementia with Lewy bodies; HS = hippocampal-sparing AD; HSc = hippocampal sclerosis; ICH = intracerebral hemorrhage; LP = limbic-predominant AD; MA = minimal atrophy AD; RID = assigned individual ID in the AD Neuroimaging Initiative dataset; SAL = subcortical arteriosclerotic leukoencephalopathy; SDH = subdural hemorrhage; TAD = typical AD; TDP-MTL = TAR DNA-binding protein in the medial temporal lobe.

severity was negatively associated with concomitance of AD and non-AD pathologies in the entorhinal cortex. This indicates that a lower value of BV:CSF showed a higher concomitance of multiple pathologies (eFigure 3, links.lww.com/WNL/C4).

Antemortem Atrophy Subtypes and Primary and Secondary Postmortem Diagnosis


The primary neuropathologic diagnosis was ADNC in all individuals (Figure 1B). Several cases had a secondary neuropathologic

Table 2 Association of Antemortem Atrophy Subtype Dimensions With AD Neuropathologic Rating Scales and Presence of Comorbid Non-AD Pathologies

Postmortem pathology	Antemortem typicality, rho (p)	Antemortem severity, rho (p)
Thal Aβ phase	-0.39 (0.035)	0.18 (0.37)
Braak Tau stage	-0.19 (0.32)	-0.18 (0.35)
Neuritic plaque	-0.40 (0.034)	0.18 (0.61)
α-Syn	-0.03 (0.86)	-0.21 (0.29)
TDP-43 inclusions	-0.49 (0.011)	-0.16 (0.46)

Abbreviations: α-syn = α-synuclein; Aβ = β-amyloid; AD = Alzheimer disease; TDP-43 = TAR DNA-binding protein 43. Overall α-syn was evaluated across brainstem-predominant, limbic/transitional, neocortical/diffuse, and amygdala-predominant stages, and overall TDP-43 was evaluated across the amygdala, hippocampus, entorhinal/inferior temporal cortex, and neocortex. α-Syn and TDP-43 pathologies were binarized to evaluate presence or absence; associations between typicality or severity and individual pathologies were evaluated using partial correlation, adjusted for field strength, age at scan, and the other dimension of subtypes (severity or typicality); rho = linear partial correlation coefficient.

Figure 2 Distribution of Postmortem AD Neuropathologies Superposed on MRI-Based Heterogeneity




Postmortem AD pathologies used to assess ADNC, encompassing the “ABC” scores of (A) Thal phase for A β , (B) Braak stage for tau, and (C) Consortium to Establish a Registry for AD neuritic plaques. All plots show antemortem MRI-based typicality on the horizontal scale, proxied by the index = $\left(\frac{\text{hippocampal volume}}{\text{cortical volume}}\right)$; all plots show antemortem MRI-based severity on the vertical scale, proxied by the Global Brain Atrophy Index = $\left(\frac{\text{total brain volume}}{\text{cerebrospinal fluid volume}}\right)$, whereby higher values correspond to lower severity. AD = Alzheimer disease; ADNC = AD neuropathologic change.

diagnosis (Figure 1C), including LB disease (n = 16, 51.610%), medial temporal TDP-43 pathology and/or hippocampal sclerosis (n = 4, 12.900%), and cerebrovascular pathology (subdural hemorrhage, intracerebral hemorrhage, and/or subcortical arteriosclerotic leukoencephalopathy [n = 3, 9.690%]). Qualitatively, cases assigned to have TDP-43 in the medial temporal lobe or hippocampal sclerosis inclined toward limbic-predominant AD or typical


AD. Cases assigned to have LB pathology tended to be limbic-predominant AD (dementia with LB pathology), hippocampal-sparing AD (amygdala-predominant LB pathology), or minimal atrophy AD (both forms). The single isolated cases with intracerebral hemorrhage, subdural hemorrhage, and subcortical arteriosclerotic leukoencephalopathy tended toward minimal atrophy AD.

Figure 3 Distribution of Postmortem Non-AD Neuropathologies Superposed on MRI-Based Heterogeneity



Postmortem non-AD pathologies including (A) α -synuclein Lewy bodies and (B) TDP-43. All plots show antemortem MRI-based typicality on the horizontal scale, proxied by the index = $\left(\frac{\text{hippocampal volume}}{\text{cortical volume}}\right)$; all plots show antemortem MRI-based severity on the vertical scale, proxied by the Global Brain Atrophy Index = $\left(\frac{\text{total brain volume}}{\text{cerebrospinal fluid volume}}\right)$, whereby higher values correspond to lower severity. A + E = TDP-43 immunoreactive inclusions are present in the amygdala and entorhinal/inferior temporal cortex; A + H + E + N = TDP-43 immunoreactive inclusions are present in the amygdala, hippocampus, entorhinal/inferior temporal cortex, and neocortex; A + H + E = TDP-43 immunoreactive inclusions are present in the amygdala, hippocampus, and entorhinal/inferior temporal cortex; AD = Alzheimer disease; TDP-43 = TAR DNA-binding protein 43.

Figure 4 Association Between Antemortem MRI-Based (A) Typicality and (B) Severity and Regional Neuropathologic Features



Associations between each of typicality or severity and presence of regional pathologies were evaluated using linear partial correlation, adjusted for field strength, age at scan, and the other dimension (severity or typicality); linear partial correlation coefficient (ρ) and significant p values are indicated. AMYG = amygdala; A β = β -amyloid (diffuse and cored plaques); CA1 = hippocampus at the level of lateral geniculate nucleus including cornu Ammonis 1 subfield; DG = hippocampus at the level of lateral geniculate nucleus including dentate gyrus; ERC = entorhinal cortex; IPL = inferior parietal lobe (angular gyrus); MFG = middle frontal gyrus; PHG = hippocampus at the level of lateral geniculate nucleus including parahippocampal gyrus; STG = superior and middle temporal gyri; Tau = phosphorylated tau assessing neurofibrillary tangles; TDP-43 = phosphorylated TAR DNA-binding protein 43 neuronal cytoplasmic inclusion; α -syn = alpha-synuclein Lewy body pathology.

The Role of Sex and APOE Status as Mediators

Corresponding to each significant association detected, we found that neither sex nor APOE status was likely a mediator of the antemortem-postmortem relationship.

Discussion

Our study investigated the relationship between antemortem atrophy subtypes and combinations of different AD and non-AD pathologies assessed postmortem. Heterogeneity in AD is a multifaceted phenomenon involving combinations of protective factors, risk factors, and concomitance of non-AD pathologies.⁷ The relative contribution of different pathologies to disease heterogeneity has been primarily reported from the postmortem (neuropathologic) perspective,^{16,24-28} with only 1 study offering an antemortem (neuroimaging) perspective,²⁹ to our knowledge. Our study serves as a direct antemortem-to-postmortem investigation examining the interplay of different pathologies in atrophy subtypes of AD.


From the antemortem perspective, we treated biological heterogeneity in atrophy as continuous phenomena,¹⁵ that is, we examined an MRI-based operationalization of the conceptual framework for AD subtypes in terms of typicality and severity.⁷ This approach is complementary to previous studies that conventionally categorize individuals into distinct subtypes.³⁰⁻³³ We observed a nonsignificant association between typicality and severity, suggesting that disease typicality (proxied by H:C) may not be influenced by disease staging or severity (proxied by BV:CSF), thus serving as orthogonal dimensions of heterogeneity. It is important, however, to note that our initial approach of

treating typicality and severity dimensions separately (while controlling for the other dimension) may be rather simplistic and deserves future exploration. This is best exemplified by cases RID 1203 and RID 1452 (Figure 1A). Despite having a lower severity (higher BV:CSF), case RID 1203 was described as hippocampal-sparing AD rather than a minimal atrophy AD. Thus, the combined contribution of typicality and severity must be factored in, that is, every individual along the typicality dimension must also be interpreted in conjunction with the corresponding severity level and vice versa.

Our key finding was that antemortem typicality, but not severity, was associated with different pathologies observed postmortem, including A β , tau, α -syn, and TDP-43. One reasoning for the lack of association between antemortem severity and postmortem pathologies could be that most individuals were at advanced disease stages (high ADNC), contributing to a low variability in postmortem disease severity. Below, we discuss the role of individual pathologies in relation to antemortem heterogeneity in atrophy.

We found an association between typicality and Thal A β stages, suggesting lower A β in the hippocampal-sparing AD atrophy subtype, which is consistent with a recent meta-analysis evaluating the proportion of A β positivity in this subtype.⁷ This result may be expected given that A β hallmark pathology in AD is rather diffuse, which may be indirectly associated with some degree of downstream atrophy.³⁴ However, we did not find a significant association of typicality with regional ratings of A β density, perhaps because A β accumulation is usually widespread and homogeneous, with little regional specificity. To some degree, this lack of regional

Figure 5 Susceptibility of Antemortem MRI-Based Heterogeneity to AD and Non-AD Neuropathologies



Associations of antemortem typicality and severity with postmortem neuropathologic features may generate the following hypotheses: (1) the orthogonal dimensions of biological heterogeneity, typicality and severity, may offer complementary information regarding the vulnerability of the brain to AD (amyloid, tau) and non-AD (α -syn, TDP-43) pathologies; and (2) limbic-predominant AD along the typicality dimension and typical AD along the severity dimension may share similar underlying biological pathway(s), which make them more susceptible to pathologies, whereas hippocampal-sparing AD along the typicality dimension and minimal AD along the severity dimension may share similar pathway(s), making them less susceptible. α -syn = α -synuclein; AD = Alzheimer disease; TDP-43 = TAR DNA-binding protein 43.

associations likely reflects the lack of topographical correspondence between $A\beta$ and atrophy because evidence suggests a closer relationship between atrophy and tau than atrophy and $A\beta$.³⁵⁻³⁷

We did not observe an association between typicality or severity and Braak NFT stages, although the AD dementia cases ($N = 26$ at Braak stage V or VI) were at relatively more advanced stages than aMCI ($N = 5$ at Braak stage III or V). This lack of association is most likely because of little variability in this measure because all but 2 cases (Braak stage III, both aMCI) were at Braak stage V or VI. When assessing NFT load regionally, however, the limbic-predominant AD atrophy subtype was associated with the presence of tau pathology in the hippocampus. This is not surprising because tau pathology is a hallmark of AD affecting the hippocampus, particularly the dentate gyrus, which is known to contain the largest density of synapses.³⁸ Thus, the presence of tau pathology may eventually be reflected in significant atrophy in the region, which is a key characteristic of the limbic-predominant AD atrophy subtype. Conversely, the hippocampal-sparing AD atrophy subtype was associated with the absence of tau

pathology in the dentate gyrus of the hippocampus. Supporting evidence for the association between atrophy and tau pathology in this subtype is not straightforward, owing to factors including the interval between assessments of these biomarkers,¹⁵ regional nonspecificity of atrophy, and disagreement of subtyping methods based on these biomarkers.³⁹ Altogether, our study is useful in providing a direct link between antemortem atrophy and postmortem tau pathology, suggesting that hippocampal atrophy relative to neocortical atrophy can track postmortem NFT subtypes.¹⁶

When assessing α -syn LB pathology regionally, we observed that the limbic-predominant AD atrophy subtype may be more prone to presence of this pathology. We also found that the parahippocampal gyrus was significantly associated with the presence of overall α -syn pathology. These findings corroborate previous postmortem neuropathologic studies showing increased α -syn pathology in typical AD and limbic-predominant AD tau subtypes.^{16,24} However, other postmortem neuropathologic studies have reported increased α -syn pathology in the hippocampal-sparing AD tau subtype.^{25,26,29} Moreover, a recent antemortem MRI study in dementia with LB observed predominance of the hippocampal-sparing atrophy subtype.⁴⁰ It must, however, be noted that most of the postmortem studies reporting the presence of α -syn pathology to date have characterized tau subtypes, which are not necessarily interchangeable with atrophy subtypes in AD.^{15,39} Therefore, future *in vivo* investigations are warranted to confirm the role of α -syn pathology in AD heterogeneity. Furthermore, α -syn LB (neocortical) pathology may potentially interact with tau (Braak stage V–VI) pathology and advanced age in our cohort, explaining atrophy in the limbic-predominant AD atrophy subtype, given that limbic atrophy is not observable in the absence of these factors.⁴¹

Our most robust findings included the association of the limbic-predominant AD atrophy subtype with the presence of TDP-43 pathology. The limbic-predominant AD tau subtype has been described to be more prone to exhibiting TDP-43 in previous postmortem studies.^{16,24,26} It is thus plausible for the limbic-predominant AD atrophy subtype to follow suit, given the topographical similarity between tau and atrophy patterns in limbic-predominant AD.¹⁵ Congruent with the report from the recent meta-analysis,⁷ our study provides an antemortem-to-postmortem validation and evidence supporting the association of the limbic-predominant AD atrophy subtype with TDP-43. We observed a gradually increasing number of brain regions being affected by TDP-43 as one moves along the typicality dimension toward limbic-predominant AD. Regional examination revealed the strongest association between typicality and presence of TDP-43 in the amygdala, an initial affected site by this pathology,²⁸ as well as in other medial temporal lobe structures (hippocampus and entorhinal cortex), shown to be affected by a recent antemortem study.⁶ As a main contributor of pathology affecting the hippocampus, TDP-43-associated hippocampal atrophy may be detectable at least 10 years before death.⁴² Thus, the limbic-predominant AD atrophy subtype is

most likely to exhibit limbic-predominant age-related TDP-43 encephalopathy neuropathologic changes.⁴³ In the absence of *in vivo* biomarkers assessing TDP-43, antemortem atrophy-based typicality (H:C) as a consistent correlate of postmortem TDP-43 in our study indicates the potential of this index as an antemortem proxy for this pathology.

Another main finding of our study was that both typicality and severity were regionally associated with concomitance of pathologies. This relationship was such that limbic-predominant AD and typical AD subtypes were associated with higher concomitance, whereas hippocampal-sparing AD and minimal atrophy AD subtypes were associated with lower concomitance. There also appears to be a region-specific effect, whereby some regions may accumulate a greater number of pathologies whereas other regions may be spared. For example, limbic-predominant AD was associated with a higher concomitance of different pathologies, particularly in the medial and superior temporal structures, and typical AD was associated with higher pathologic concomitance in the entorhinal cortex. Of interest, hippocampal structures including the dentate gyrus and CA1 demonstrated a generally lower concomitance than other regions. The divergent reports mentioned previously on α -syn pathology being associated with limbic-predominant and typical atrophy subtypes may be due to the higher susceptibility of the subtypes to multiple or mixed pathologies.

Finally, although qualitative, individual-level secondary postmortem diagnoses aided in providing greater confidence to our quantitative findings. Two cases with lower H:C index (toward limbic-predominant AD) were diagnosed to have TDP-43 in the medial temporal region, consistent with our main quantitative findings. Two additional cases with lower H:C index were diagnosed to have hippocampal sclerosis, which is known to correlate well with TDP-43 pathology.^{16,24} Five of 6 cases with relatively higher H:C index (toward hippocampal-sparing AD) were assigned to have amygdala-predominant LB pathology, a distinct pathologic entity.⁴ Whether/how the presence of LB pathology in the amygdala plays a role in the disposition of the hippocampal-sparing AD atrophy subtype to the pathology remains to be seen. Three cases with relatively higher H:C (toward hippocampal-sparing AD) and higher BV:CSF (toward minimal atrophy AD) indices were diagnosed with cerebrovascular pathologies. Although the lack of variability in the measure of cerebrovascular disease did not allow us to account for it in our quantitative analyses, these qualitative observations align with recent evidence, showing that cerebrovascular disease may particularly affect hippocampal-sparing AD⁴⁴ and minimal atrophy AD subtypes.^{44,45}

Considering our current findings, we propose 2 hypotheses for future work, as larger antemortem-postmortem datasets become available (Figure 5): (1) biological heterogeneity, characterized by the orthogonal dimensions of typicality and severity, captures different aspects of vulnerability of the brain to AD and non-AD pathologies. Although typicality may be relatively more sensitive to individual pathologies varying

regionally, severity may predominantly reflect a cumulative contribution of several pathologies, measured as concomitance; (2) limbic-predominant AD and typical AD subtypes may follow a unique biological pathway that tends to be affected by greater accumulation, interaction, and concomitance of various pathologies, distinct from the pathway followed by the hippocampal-sparing AD subtype that may be less affected. It is unclear which pathway the minimal atrophy AD subtype may follow: at antemortem, individuals tending toward minimal atrophy AD were at early disease stages (i.e., amnesic mild cognitive impairment) and could have eventually progressed into one of the other 3 subtypes, thus possibly following either of the 2 hypothesized pathways; at postmortem, however, many of these individuals showed high ADNC despite having minimal atrophy, suggesting that minimal atrophy AD may share the pathway common to the hippocampal-sparing AD subtype of being less affected by concomitance of various pathologies. Although our current and recent works^{46,47} provide initial support, these hypotheses need to be tested by future studies to understand their potential validity across different modalities (heterogeneity assessed by measures other than atrophy), pathologies (e.g., vascular burden), and disease stages (including predementia cases).

Our study has some limitations. First, the sample size of our cohort was limited, which may reduce the power to detect significant associations and generalize findings. However, our sample size was comparable with previous studies combining antemortem and postmortem data.^{48,49} Despite the size, we observed representation of 4 subtypes, and we chose methodologies proportionate to this limited sample size by modeling heterogeneity with continuous measures (typicality and severity) and analyzing heterogeneity using partial correlation models to maximize statistical power. Second, postmortem pathologies were only available as semiquantitative scores (i.e., gross burden of pathology), which we further binarized for the presence/absence of pathologies for sufficient statistical power. These scales may not be as sensitive as quantitative scores obtained from digital histology techniques (e.g., specific counts, density, or percentage of pathology per region). Third, most of the individuals showed a high ADNC (low variability in postmortem severity), which may have influenced the finding that the associations of antemortem MRI typicality with postmortem pathologies were stronger than those of MRI severity. Future investigations should include a broader range of pathologic severity to fully explore associations for the severity dimension. Finally, all data were sourced from the ADNI, known to have relatively strict inclusion criteria. Therefore, our current findings would need to be further validated by future studies using less restrictive and more heterogeneous cohorts.

In conclusion, we examined the relationship between antemortem MRI-based atrophy subtypes (modeled as continuous phenomena) and postmortem neuropathology in AD. In our cohort, antemortem typicality shared a stronger overall and region-specific association with different postmortem pathologies, including A β , tau, α -synuclein, and TDP-43,

compared with antemortem severity. This suggests that the novel operationalization of biological heterogeneity in AD including typicality as a continuum is a promising proxy for the presence and regional distribution of pathologies, irrespective of disease staging (severity). Thus, factoring in contributions of core AD and comorbid non-AD pathologies toward biological heterogeneity in unspecific markers of neurodegeneration may subsequently serve as an avenue for precision medicine and future multifactorial therapies.

Study Funding

This study was funded by the Swedish Foundation for Strategic Research (SSF); the Strategic Research Programme in Neuroscience at Karolinska Institutet (StratNeuro); the Swedish Research Council (VR); the regional agreement on medical training and clinical research (ALF) between Stockholm County Council and Karolinska Institutet; Center for Innovative Medicine (CIMED); the Swedish Alzheimer Foundation; the Swedish Brain Foundation; the Åke Wiberg Foundation; Demensfonden; Stiftelsen Olle Engkvist Byggmästare; Birgitta och Sten Westerberg; Foundation for Geriatric Diseases at Karolinska Institutet; Loo och Hans Ostermans stiftelse för medicinsk forskning; Stiftelsen För Gamla Tjänarinnor; and Gun and Bertil Stohne Stiftelse. M.J.G. is supported by the “Miguel Servet” program [CP19/00031] and a research grant [PI20/00613] of the Instituto de Salud Carlos III-Fondo Europeo de Desarrollo Regional (ISCIII-FEDER). The funding sources did not have any involvement in the study design; collection, analysis, and interpretation of data; writing of the report; and the decision to submit the article for publication. Data collection and sharing for this study was funded by the Alzheimer’s Disease Neuroimaging Initiative (ADNI) (NIH Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). The ADNI is funded by the National Institute on Aging, by the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: Alzheimer’s Association; Alzheimer’s Drug Discovery Foundation; BioClinica, Inc.; Biogen Idec Inc.; Bristol-Myers Squibb Company; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; GE Healthcare; Innogenetics, N.V.; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC; Johnson & Johnson Pharmaceutical Research & Development LLC; Medpace, Inc.; Merck & Co., Inc.; Meso Scale Diagnostics, LLC; NeuroRx Research; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Synarc Inc.; and Takeda Pharmaceutical Company. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the NIH (fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer’s Disease Cooperative Study at the University of California, San Diego. ADNI data are disseminated by the Laboratory of Neuro Imaging at the University of California, Los Angeles.

Disclosures

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org/N](https://doi.org/10.1101/2021.09.06.21263162) for full disclosures.

Publication History

Previously published at medRxiv: <https://doi.org/10.1101/2021.09.06.21263162>. Received by *Neurology* October 22, 2021. Accepted in final form March 4, 2022. Submitted and externally peer reviewed. The handling editors were Rawan Tarawneh, MD, and Brad Worrall, MD, MSc.

Appendix Authors

Name	Location	Contribution
Rosaleena Mohanty, PhD	Division of Clinical Geriatrics, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Stockholm, Sweden	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; and analysis or interpretation of data
Daniel Ferreira, PhD	Division of Clinical Geriatrics, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Stockholm, Sweden; Department of Radiology, Mayo Clinic, Rochester, MN	Drafting/revision of the manuscript for content, including medical writing for content, and study concept or design
Simon Frerich, MSc	Division of Clinical Geriatrics, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Stockholm, Sweden; Institute for Stroke and Dementia Research, University Hospital, Ludwig-Maximilian-University (LMU) Munich, Germany	Analysis or interpretation of data
J-Sebastian Muehlboeck, MSc	Division of Clinical Geriatrics, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Stockholm, Sweden	Analysis or interpretation of data
Michel J. Grothe, PhD	Unidad de Trastornos del Movimiento, Servicio de Neurología y Neurofisiología Clínica, Instituto de Biomedicina de Sevilla, Hospital Universitario Virgen del Rocío/CSIC/ Universidad de Sevilla, Spain; Clinical Dementia Research Section, German Center for Neurodegenerative Diseases (DZNE), Rostock, Germany	Drafting/revision of the manuscript for content, including medical writing for content, and study concept or design
Eric Westman, PhD	Division of Clinical Geriatrics, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Stockholm, Sweden; Department of Neuroimaging, Centre for Neuroimaging Sciences, Institute of Psychiatry, Psychology and Neuroscience, King’s College London, UK	Drafting/revision of the manuscript for content, including medical writing for content; study concept or design; analysis or interpretation of data; and other

References

1. Barker WW, Luis CA, Kashuba A, et al. Relative frequencies of Alzheimer disease, Lewy body, vascular and frontotemporal dementia, and hippocampal sclerosis in the State of Florida Brain Bank. *Alzheimer Dis Assoc Disord*. 2002;16(4):203-212.
2. Rahimi J, Kovacs GG. Prevalence of mixed pathologies in the aging brain. *Alzheimer's Res Ther*. 2014;6(9):1-11.
3. Spire-Jones TL, Attems J, Thal DR. Interactions of pathological proteins in neurodegenerative diseases. *Acta Neuropathol*. 2017;134(2):187-205.
4. Uchikado H, Lin WL, DeLucia MW, Dickson DW. Alzheimer disease with amygdala Lewy bodies: a distinct form of α -synucleinopathy. *J Neuropathol Exp Neurol Am Assoc Neuropathologists*. 2006;65(7):685-697.
5. Josephs KA, Murray ME, Whitwell JL, et al. Updated TDP-43 in Alzheimer's disease staging scheme. *Acta Neuropathol*. 2016;131(4):S71-S85.
6. de Flores R, Wisse LEM, Das SR, et al. Contribution of mixed pathology to medial temporal lobe atrophy in Alzheimer's disease. *Alzheimer's Dement*. 2020;16(6):843-852.
7. Ferreira D, Nordberg A, Westman E. Biological subtypes of Alzheimer disease: a systematic review and meta-analysis. *Neurol AAN Enterprises*. 2020;94(10):436-448.
8. Montine TJ, Phelps CH, Beach TG, et al. National Institute on Aging-Alzheimer's Association guidelines for the neuropathologic assessment of Alzheimer's disease: a practical approach. *Acta Neuropathol*. 2012;123(1):1-11.
9. Muehlboeck J, Westman E, Simmons A. TheHiveDB image data management and analysis framework. *Front Neuroinform Front*. 2014;7:49.
10. Desikan RS, Ségonne F, Fischl B, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage*. 2006;31(3):968-980.
11. Fischl B, Salat DH, Busa E, et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron*. 2002;33(3):341-355.
12. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12(3):189-198.
13. Crane PK, Carle A, Gibbons LE, et al. Development and assessment of a composite score for memory in the Alzheimer's Disease Neuroimaging Initiative (ADNI). *Brain Imaging Behav*. 2012;6(4):502-516.
14. Gibbons LE, Carle AC, Mackin RS, et al. A composite score for executive functioning, validated in Alzheimer's Disease Neuroimaging Initiative (ADNI) participants with baseline mild cognitive impairment. *Brain Imaging Behav*. 2012;6(4):517-527.
15. Mohanty R, Ferreira D, Nordberg A, Westman E. Associations between different tau-PET patterns and longitudinal atrophy in the Alzheimer's disease continuum. *medRxiv*. 2021.
16. Murray ME, Graff-Radford NR, Ross OA, Petersen RC, Duara R, Dickson DW. Neuropathologically defined subtypes of Alzheimer's disease with distinct clinical characteristics: a retrospective study. *Lancet Neurol*. 2011;10(9):785-796.
17. Orellana C, Ferreira D, Muehlboeck J-S, et al. Measuring global brain atrophy with the brain volume/cerebrospinal fluid index: normative values, cut-offs and clinical associations. *Neurodegener Dis*. 2016;16(1-2):77-86.
18. Franklin EE, Perrin RJ, Vincent B, et al. Brain collection, standardized neuropathologic assessment, and comorbidity in Alzheimer's Disease Neuroimaging Initiative 2 participants. *Alzheimer's Dement*. 2015;11(7):815-822.
19. McKeith IG, Dickson DW, Lowe J, et al. Diagnosis and management of dementia with Lewy bodies: third report of the DLB Consortium. *Neurology*. 2005;65(12):1863-1872.
20. Katsumata Y, Fardo DW, Kukull WA, Nelson PT. Dichotomous scoring of TDP-43 proteinopathy from specific brain regions in 27 academic research centers: associations with Alzheimer's disease and cerebrovascular disease pathologies. *Acta Neuropathol Commun Biomed Cent*. 2018;6(1):1-11.
21. Wisse LEM, de Flores R, Xie L, et al. Pathological drivers of neurodegeneration in suspected non-Alzheimer's disease pathophysiology. *Alzheimer's Res Ther*. 2021;13(1):1-11.
22. Spina S, La Joie R, Petersen C, et al. Comorbid neuropathological diagnoses in early versus late-onset Alzheimer's disease. *Brain*. 2021;144(7):2186-2198.
23. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol*. 1986;51(6):1173.
24. Janocko NJ, Brodersen KA, Soto-Ortolaza AI, et al. Neuropathologically defined subtypes of Alzheimer's disease differ significantly from neurofibrillary tangle-predominant dementia. *Acta Neuropathol*. 2012;124(5):681-692.
25. Jellinger KA. Neuropathological subtypes of Alzheimer's disease. *Acta Neuropathol*. 2012;123(1):153.
26. Josephs KA, Whitwell JL, Tosakulwong N, et al. TAR DNA-binding protein 43 and pathological subtype of Alzheimer's disease impact clinical features. *Ann Neurol*. 2015;78(5):697-709.
27. Murray ME, Cannon A, Graff-Radford NR, et al. Differential clinicopathologic and genetic features of late-onset amnesic dementias. *Acta Neuropathol*. 2014;128(3):411-421.
28. Hu WT, Josephs KA, Knopman DS, et al. Temporal lobar predominance of TDP-43 neuronal cytoplasmic inclusions in Alzheimer disease. *Acta Neuropathol*. 2008;116(2):215.
29. Whitwell JL, Dickson DW, Murray ME, et al. Neuroimaging correlates of pathologically defined subtypes of Alzheimer's disease: a case-control study. *Lancet Neurol*. 2012;11(10):868-877.
30. Ferreira D, Verhagen C, Hernández-Cabrera JA, et al. Distinct subtypes of Alzheimer's disease based on patterns of brain atrophy: longitudinal trajectories and clinical applications. *Sci Rep*. 2017;7:46263.
31. Poulakis K, Pereira JB, Mecocci P, et al. Heterogeneous patterns of brain atrophy in Alzheimer's disease. *Neurobiol Aging*. 2018;65:98-108.
32. Byun MS, Kim SE, Park J, et al. Heterogeneity of regional brain atrophy patterns associated with distinct progression rates in Alzheimer's disease. *PLoS One*. 2015;10(11):e0142756.
33. Risacher SL, Anderson WH, Charil A, et al. Alzheimer disease brain atrophy subtypes are associated with cognition and rate of decline. *Neurology*. 2017;89(21):2176-2186.
34. Iaccarino L, Tammewar G, Ayakta N, et al. Local and distant relationships between amyloid, tau and neurodegeneration in Alzheimer's Disease. *Neuroimage Clin*. 2018;17:452-464.
35. Nelson PT, Alafuzoff I, Bigio EH, et al. Correlation of Alzheimer disease neuropathologic changes with cognitive status: a review of the literature. *J Neuropathol Exp Neurol*. 2012;71(5):362-381.
36. Gordon BA, McCullough A, Mishra S, et al. Cross-sectional and longitudinal atrophy is preferentially associated with tau rather than amyloid β positron emission tomography pathology. *Alzheimer's Dement*. 2018;10:245-252.
37. Ossenkoppele R, Smith R, Mattsson-Carlsson N, et al. Accuracy of tau positron emission tomography as a prognostic marker in preclinical and prodromal Alzheimer disease: a head-to-head comparison against amyloid positron emission tomography and magnetic resonance imaging. *JAMA Neurol*. 2021;78(8):961-971.
38. Lace G, Savva GM, Forster G, et al. Hippocampal tau pathology is related to neuroanatomical connections: an ageing population-based study. *Brain*. 2009;132(pt 5):1324-1334.
39. Mohanty R, Mårtensson G, Poulakis K, et al. Comparison of subtyping methods for neuroimaging studies in Alzheimer's disease: a call for harmonization. *Brain Commun*. 2020;2(2):fcaa192.
40. Oppedal K, Ferreira D, Cavallin L, et al. A signature pattern of cortical atrophy in dementia with Lewy bodies: a study on 333 patients from the European DLB consortium. *Alzheimer's Dement*. 2019;15(3):400-409.
41. Ferman TJ, Aoki N, Boeve BF, et al. Subtypes of dementia with Lewy bodies are associated with α -synuclein and tau distribution. *Neurology*. 2020;95(2):e155-e165.
42. Josephs KA, Dickson DW, Tosakulwong N, et al. Rates of hippocampal atrophy and presence of post-mortem TDP-43 in patients with Alzheimer's disease: a longitudinal retrospective study. *Lancet Neurol*. 2017;16(11):917-924.
43. Nelson PT, Dickson DW, Trojanowski JQ, et al. Limbic-predominant age-related TDP-43 encephalopathy (LATE): consensus working group report. *Brain*. 2019;142(6):1503-1527.
44. Ferreira D, Shams S, Cavallin L, et al. The contribution of small vessel disease to subtypes of Alzheimer's disease: a study on cerebrospinal fluid and imaging biomarkers. *Neurobiol Aging*. 2018;70:18-29.
45. Ferreira D, Wahlund L-O, Westman E. The heterogeneity within Alzheimer's disease. *Aging (Albany NY)*. *Impact Journals*. 2018;10(11):3058.
46. Poulakis K, Pereira JB, Muehlboeck J, et al. Stage vs. Subtype Hypothesis in Alzheimer's Disease: A Multi-Cohort and Longitudinal Bayesian Clustering Study. March 4, 2021. Accessed May 30, 2022. <http://dx.doi.org/10.2139/ssrn.3797614>.
47. Ferreira D, Mohanty R, Murray M, Nordberg A, Kantarci K, Westman E. Does a truly hippocampal sparing subtype of Alzheimer's disease really exist? 2021. *Alzheimer's Assoc Int Conf*. 2021;17(S4):e056368.
48. Teipel SJ, Fritz H-C, Grothe MJ, Initiative ADN. Neuropathologic features associated with basal forebrain atrophy in Alzheimer disease. *Neurol*. 2020;95(10):e1301-e1311.
49. Grothe MJ, Moscoso A, Ashton NJ, et al. Associations of fully automated CSF and novel plasma biomarkers with Alzheimer disease neuropathology at autopsy. *Neurology*. 2021;97(12):e1229-e1242.

Neurology[®]

Neuropathologic Features of Antemortem Atrophy-Based Subtypes of Alzheimer Disease

Rosaleena Mohanty, Daniel Ferreira, Simon Frerich, et al.
Neurology 2022;99:e323-e333 Published Online before print May 24, 2022
DOI 10.1212/WNL.0000000000200573

This information is current as of May 24, 2022

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/99/4/e323.full
References	This article cites 47 articles, 5 of which you can access for free at: http://n.neurology.org/content/99/4/e323.full#ref-list-1
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/99/4/e323.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Alzheimer's disease http://n.neurology.org/cgi/collection/alzheimers_disease MRI http://n.neurology.org/cgi/collection/mri
Errata	An erratum has been published regarding this article. Please see next page or: /content/100/3/165.1.full.pdf
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.



Disputes & Debates: Editors' Choice

Steven Galetta, MD, FAAN, Editor
Aravind Ganesh, MD, DPhil, FRCPC, Deputy Editor
Ariane Lewis, MD, Deputy Editor
James E. Siegler III, MD, Deputy Editor

Editors' Note: Benefits and Risks of Epilepsy Surgery in Patients With Focal Cortical Dysplasia Type 2 in the Central Region

In the research article entitled "Benefits and Risks of Epilepsy Surgery in Patients With Focal Cortical Dysplasia Type 2 in the Central Region," Chassoux et al. described their findings from a retrospective review of preoperative and postoperative data for 60 patients with focal cortical dysplasia type 2 (FCD2) in the central region who underwent surgical resection. They reported that surgical resection led to seizure freedom in 88% of patients, and although 87% had early transitory postoperative deficits, 40% of these patients had total recovery. Abe noted that patients with FCD2 can experience neurologic deficits due to the location of the dysplasia and asked the authors to describe preoperative neurologic deficits for patients included in this study. Chassoux reported that 18% of patients experienced a preoperative deficit that was either permanent or fluctuating and they all experienced transitory worsening of this deficit postoperatively, but 36% subsequently had functional improvement. Chassoux also noted that 84% of patients with a normal neurologic examination preoperatively experienced a postoperative deficit.

Ariane Lewis, MD, and Steven Galetta, MD, FAAN
Neurology® 2023;100:162. doi:10.1212/WNL.0000000000206737

Reader Response: Benefits and Risks of Epilepsy Surgery in Patients With Focal Cortical Dysplasia Type 2 in the Central Region

Kazuo Abe (Osaka, Japan)
Neurology® 2023;100:162. doi:10.1212/WNL.0000000000206738

I read the article by Chassoux et al.¹ with interest, and I have some questions that may be outside the focus of their study. The authors described that resections were associated with minor/moderate deficits and total recovery was observed in 40% of patients. Did any of these patients have neurologic deficits before resections? For example, the patient demonstrated in the MRI images showed focal cortical dysplasia near the precentral knob. Injury to this area has been reported to show isolated motor palsy in the hand, mimicking a peripheral nerve palsy.^{2,3} Can the authors please comment on the neurologic findings of these patients?

1. Chassoux F, Mellerio C, Laurent A, Landre E, Turak B, Devaux B. Benefits and risks of epilepsy surgery in patients with focal cortical dysplasia type 2 in the central region. *Neurology*. 2022;99:e11-e22.
2. Kim JS. Predominant involvement of a particular group of fingers due to small, cortical infarction. *Neurology*. 2001;56:1677-1682.
3. Komatsu K, Fukutake T, Hattori T. Isolated shoulder paresis caused by a small cortical infarction. *Neurology*. 2003;61:1457.

Copyright © 2023 American Academy of Neurology

Author Response: Benefits and Risks of Epilepsy Surgery in Patients With Focal Cortical Dysplasia Type 2 in the Central Region

Francine Chassoux (Paris)

Neurology® 2023;100:163. doi:10.1212/WNL.0000000000206739

We are grateful for the opportunity to respond to the comments by Dr. Abe on our research.¹ As described in the Results section of the article, 11 patients had a preoperative deficit that was either permanent (including a spastic component in 2 and arm atrophy in 1) or fluctuating, according to the seizure frequency (clumsiness or hemineglect). Postoperatively, all of them had a transitory worsening (which was major in 5 patients), and then, a recovery included functional improvement in 4. Among the 49 patients with normal preoperative examination, 41 had a postoperative deficit (which was major in 14).

Whatever the preoperative status, major deficits only occurred after resections of the primary motor cortex (PMC) or supplementary motor area (SMA). PMC resections were followed by pure motor deficits involving a limb or a part of limb with a central topography. Resections performed near the hand knob induced a brachial deficit predominant on the hand, but without selective involvement of fingers. SMA resections were followed by massive hemiplegia with proximal predominance and speech disturbances. A total or subtotal recovery occurred in most patients. Minor to moderate deficits included mild hand clumsiness, slow finger tapping, or a limitation for carrying heavy loads. Permanent major deficits occurred in 3 patients, who presented with central characteristics including spasticity. Contrasting with the observations based on small cortical infarctions, none of the postoperative motor deficits mimicked a peripheral nerve palsy.

1. Chassoux F, Mellerio C, Laurent A, Landre E, Turak B, Devaux B. Benefits and risks of epilepsy surgery in patients with focal cortical dysplasia type 2 in the central region. *Neurology*. 2022;99(1):e11-e22. doi:10.1212/WNL.0000000000200345.

Copyright © 2023 American Academy of Neurology

Editors' Note: Association of Diabetes and Hypertension With Brain Structural Integrity and Cognition in the Boston Puerto Rican Health Study Cohort

In the research article entitled “Association of Diabetes and Hypertension With Brain Structural Integrity and Cognition in the Boston Puerto Rican Health Study Cohort,” Guan et al. reported that comorbid diabetes and hypertension is associated with a greater degree of brain structural disruptions than hypertension alone. Kawada commented that comorbid diabetes and hypertension was not associated with a greater degree of structural or functional impairment than diabetes or hypertension alone in a study by Newby et al. Koo and Guan responded that the relationship between diabetes and hypertension and brain structural integrity and cognition may vary across cohorts due to modification by social determinants of health. Kawada also noted that there is a bidirectional relationship between diabetes and dementia that should be considered when interpreting results of the study conducted by Guan et al., to which Koo and Guan replied that while they do not have data on β -amyloid and dementia for this cohort, this could be a future research direction.

Ariane Lewis, MD, and Steven Galetta, MD, FAAN

Neurology® 2023;100:163. doi:10.1212/WNL.0000000000206740

Author disclosures are available upon request (journal@neurology.org).

Reader Response: Association of Diabetes and Hypertension With Brain Structural Integrity and Cognition in the Boston Puerto Rican Health Study Cohort

Tomoyuki Kawada (Tokyo)

Neurology® 2023;100:164. doi:10.1212/WNL.0000000000206741

The research by Guan et al.¹ reports that older patients with type 2 diabetes (T2D) and hypertension (HTN) comorbidity presented with increased brain structural change and advanced cognitive impairment. I have a few comments.

First, Newby et al.² reported the increased risk of brain damage and cognitive impairment in patients with T2D and HTN. The adverse change in the pallidum was predominant in patients with T2D and HTN comorbidity. By contrast, total gray volume and verbal-numerical reasoning were disturbed in patients with only T2D, and the symbol digit substitution task was disturbed in patients with only HTN. Comorbidity of T2D and HTN did not always present greater brain damage and functional impairment than in participants with HTN or T2D alone.

Second, Gottesman et al.³ evaluated metabolic risk and amyloid β (A β) burden with subsequent dementia. The adjusted hazard ratios of midlife HTN with late life T2D and A β deposition for dementia incidence significantly increased. Shigemori et al.⁴ reported that A β was found to be secreted from β -cells of the pancreas, along with insulin, on glucose stimulation. A bidirectional relationship between T2D and dementia should be considered for the analysis.

1. Guan Y, Ebrahimzadeh SA, Cheng CH, et al. Association of diabetes and hypertension with brain structural integrity and cognition in the Boston Puerto Rican Health Study cohort. *Neurology*. 2022;98(15):e1534-e1544.
2. Newby D, Garfield V. Understanding the inter-relationships of type 2 diabetes and hypertension with brain and cognitive health: a UK Biobank study. *Diabetes Obes Metab*. 2022;24(5):938-947.
3. Gottesman RF, Wu A, Coresh J, et al. Associations of vascular risk and amyloid burden with subsequent dementia. *Ann Neurol*. 2022;92(4):607-619.
4. Shigemori K, Nomura S, Umeda T, Takeda S, Tomiyama T. Peripheral A β acts as a negative modulator of insulin secretion. *Proc Natl Acad Sci USA*. 2022;119(12):e2117723119.

Copyright © 2023 American Academy of Neurology

Author Response: Association of Diabetes and Hypertension With Brain Structural Integrity and Cognition in the Boston Puerto Rican Health Study Cohort

Bang-Bon Koo (Boston) and Yi Guan (Boston)

Neurology® 2023;100:164–165. doi:10.1212/WNL.0000000000206742

We appreciate the comment on our article.¹ We agree that participants who presented with type 2 diabetes (T2D) or hypertension (HTN) alone may show unique brain structural changes compared with those with both comorbidities. The reader's conclusion that worse brain and cognitive outcomes in comorbid individuals is consistent with our findings. Moreover, we wish to emphasize 2 points.

First, the effects of T2D and HTN vary significantly between different cohorts. The outcome of T2D can be influenced by race, ethnicity, and social factors.² In the study conducted by Newby and Garfield,³ the UK Biobank population consists of primarily non-Hispanic White individuals, whereas our Boston Puerto Rican Health Study cohort consists of all Hispanic individuals. Second, we are also interested in studying the effects of T2D on brain imaging measures, but we were limited by sample size, as mentioned in our article. We are collecting more MRI scans and will apply our analytical framework to a larger sample in the future. Studying the association between T2D, dementia, and underlying mechanisms related to A β will be valuable for

Author disclosures are available upon request (journal@neurology.org).

integrating cardiovascular risk factors into dementia biomarker research. Our current sample does not contain information regarding A β and dementia diagnosis, but we will consider your suggestion in the future.

1. Guan Y, Ebrahimzadeh SA, Cheng CH, et al. Association of diabetes and hypertension with brain structural integrity and cognition in the Boston Puerto Rican Health Study cohort. *Neurology*. 2022;98(15):e1534-e1544.
2. Walker RJ, Strom Williams J, Egede LE. Influence of race, ethnicity and social determinants of health on diabetes outcomes. *Am J Med Sci*. 2016;351(4):366-373.
3. Newby D, Garfield V. Understanding the inter-relationships of type 2 diabetes and hypertension with brain and cognitive health: a UK Biobank study. *Diabetes Obes Metab*. 2022;24(5):938-947.

Copyright © 2023 American Academy of Neurology

CORRECTIONS

Neuropathologic Features of Antemortem Atrophy-Based Subtypes of Alzheimer Disease

Neurology® 2022;100:165. doi:10.1212/WNL.000000000201298

In the Research Article “Neuropathologic Features of Antemortem Atrophy-Based Subtypes of Alzheimer Disease” by Mohanty et al.,¹ the abbreviation for “Mini-Mental State Examination” should read “MMSE.” The authors regret the error.

Reference

1. Mohanty R, Ferreira D, Frerich S, et al. Neuropathologic features of antemortem atrophy-based subtypes of Alzheimer disease. *Neurology*. 2022;99:e323-e333.

Association of Lower Extremity Peripheral Nerve Impairment and the Risk of Dementia

Bringing the Peripheral Nervous System Closer to Center

Neurology® 2022;100:165. doi:10.1212/WNL.000000000201224

In the Editorial “Association of Lower Extremity Peripheral Nerve Impairment and the Risk of Dementia: Bringing the Peripheral Nervous System Closer to Center” by Shuman Paretsky et al.,¹ Dr. Shuman Paretsky’s degree should appear as “PhD.” The authors regret the error.

Reference

1. Shuman Paretsky M, Roman G. Association of lower extremity peripheral nerve impairment and the risk of dementia: bringing the peripheral nervous system closer to center. *Neurology*. 2022;98:741-742.