to-point representation of each finger to neurons located in the motor cortex in a way that the radial finger area is located laterally and the ulnar finger area medially,5 recent authors advocate overlapping representation or group activation theories.⁶ Isolated weakness of a single finger secondary to a motor cortical lesion observed in our patient seems to agree more with the former theory. Unexpectedly, the MRI lesion was located slightly medial to the center of the hand knob (see the figure), which does not agree with a previous report that predominant weakness of radialsided fingers is usually caused by laterally located infarcts whereas weakness of ulnar fingers is related to medial lesions in the precentral knob. Perhaps, there may be considerable individual variation of motor topography, which also was observed in the

Regarding the pathogenesis of stroke, our patient had atrial fibrillation, mobile aortic atheroma, and normal MR angiogram findings except for the posterior cerebral artery occlusion. Moreover, two infarcts developed in succession in different vascular territories. Therefore, an embolism is a likely pathogenic mechanism. This seems to be consistent with the previous observation that paresis of the radial-sided fingers was more often associated with embolic mechanisms as compared with ulnar-sided finger weakness.7

From the Department of Neurology, University of Ulsan, Asan Medical Center. Seoul. Korea.

Received September 2, 2001. Accepted in final form November 29, 2001.

Address correspondence and reprint requests to Dr. Jong S. Kim, Department of Neurology, Asan Medical Center, Song-Pa PO Box 145, Seoul 138-600, South Korea; e-mail: jongskim@www.amc.seoul.kr


Copyright © 2002 by AAN Enterprises, Inc.


References

- 1. Terao Y, Hayashi H. Kanda T, Tanabe H. Discrete cortical infarction with prominent impairment of thumb flexion. Stroke 1993;24:2118-
- Lee P-H, Han S-W, Heo JH. Isolated weakness of the fingers in cortical infarction. Neurology 1998;50:823-824.
- Phan TG, Evans BA, Huston J. Pseudoulnar palsy from a small infarct of the precentral knob. Neurology 2000;54:2185.
- Yousry TA, Schmid UD, Alkadhi H, et al. Localization of the motor and hand area to a knob on the precentral gyrus. Brain 1997;120:141-157.
- Penfield W, Boldrey E. Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. Brain 1937; 60:389-443.
- 6. Sanes JN, Donoghue JP. Static and dynamic organization of motor cortex. In: Freund H-J, Sabel BA, Witte W, eds. Brain plasticity, advances in neurology, Vol. 73. Philadelphia: Lippincott-Raven, 1997;277-296.
- 7. Kim JS. Predominant involvement of a particular group of fingers due to small, cortical infarction. Neurology 2001;56:1677-1682.

Neuro *Images*










Figure. Cerebral multislice CT angiography (MS-CTA): normal peripheral anterior cerebral artery (ACA) branches on the right side (arrows), hypoperfusion of corresponding contralateral branches (A). Cervical MS-CTA: left internal carotid artery stenosis (B) with clarification (arrowheads) and intraluminal thrombus (dotted arrow); endoluminal view shows apex of the thrombus (C). (D) Noncontrast head CT: cortical ACA infarction affecting the paramedian front and the parietal lobe (arrows).

Intraluminal carotid thrombus

Markus Busch, MD, Randolf Klingebiel, MD, Georg Bohner, MD, Florian Masuhr, MD, Berlin, Germany

A 61-year-old hypertensive man presented 1 hour after sudden onset of right-sided hemiparesis and hemineglect. Noncontrast head CT was unremarkable, but multislice CT angiography (MS-CTA) demonstrated left anterior cerebral artery (ACA) branch occlusion and ipsilateral internal carotid artery (ICA) stenosis with an intraluminal thrombus (figure, A through C). Consequently, systemic thrombolysis with recombinant tissue plasminogen activator was initiated. The next day, head CT depicted a small cortical ACA infarction (figure, D).

Isolated ACA infarction is rare and is most commonly caused by artery-to-artery embolism from atherothrombotic ICA stenosis.^{1,2} MS-CTA is a new high-resolution imaging modality that allows a quick, concurrent, and minimally invasive visualization of cervical and cerebral vessels, including virtual endovascular views.

^{1.} Gacs G, Fox AJ, Barnett HJ, Vinuela F. Occurrence and mechanisms of occlusion of the anterior cerebral artery. Stroke 1983;14:952-959

Bogousslavsky J, Regli F. Anterior cerebral artery territory infarction in the Lausanne Stroke Registry. Arch Neurol 1990;47:144-150.



Intraluminal carotid thrombus

Markus Busch, Randolf Klingebiel, Georg Bohner, et al. Neurology 2002;58;986 DOI 10.1212/WNL.58.6.986

This information is current as of March 26, 2002

Updated Information & including high resolution figures, can be found at:

Services http://n.neurology.org/content/58/6/986.full

References This article cites 2 articles, 1 of which you can access for free at:

http://n.neurology.org/content/58/6/986.full#ref-list-1

Subspecialty Collections This article, along with others on similar topics, appears in the

following collection(s):

All Cerebrovascular disease/Stroke

http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke

All Imaging

http://n.neurology.org/cgi/collection/all_imaging

CT

http://n.neurology.org/cgi/collection/ct

Embolism

http://n.neurology.org/cgi/collection/embolism

Permissions & Licensing Information about reproducing this article in parts (figures, tables) or in

its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints Information about ordering reprints can be found online:

http://n.neurology.org/subscribers/advertise

Neurology ® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

