



Figure. (A) Diffusion-weighted imaging performed on day 1 of symptoms shows no evidence of acute ischemia. (B) Repeat study performed 2 days later reveals a left lateral medullary infarction.

False-negative diffusion-weighted imaging with lateral medullary infarction

Rakesh Khatri, MD; James Leach, MD; and Matthew L. Flaherty, MD

A 55-year-old¹ man awoke with left ear pain, vertigo, vomiting, left facial numbness, and ataxic gait. MRI including diffusion-weighted imaging (DWI) performed 2 hours later showed no evidence of acute infarction (figure, A). Repeat MRI 2 days later showed DWI and FLAIR changes consistent with a left lateral medullary infarction (figure, B).

False-negative DWI has been reported in 5% of ischemic stroke cases, most commonly brainstem infarcts imaged within 24 hours of onset. Possible explanations include lesions too small for the resolution of the DWI echoplanar sequence, insufficient signal-to-noise ratio in the first hours after onset, and magnetic susceptibility artifacts causing image

distortions.¹ The clinical history and examination remain fundamental aspects of patient assessment in the era of advanced neuroimaging.

Reference

1. Oppenheim C, Stanescu R, Dormont D, et al. False-negative diffusion-weighted MR findings in acute ischemic stroke. *AJNR Am J Neuroradiol* 2000;21:1434-1440.

From the Departments of Neurology (R.K., M.L.F.) and Radiology (J.L.), University of Cincinnati College of Medicine, Cincinnati, OH.

Disclosure: The authors report no conflicts of interest.

Address correspondence and reprint requests to Dr. Matthew L. Flaherty, 231 Albert Sabin Way, MSB Room 5161B, University of Cincinnati Medical Center, Cincinnati, OH 45267-0525; e-mail: matthew.flaherty@uc.edu

Neurology[®]

False-negative diffusion-weighted imaging with lateral medullary infarction

Rakesh Khatri, James Leach and Matthew L. Flaherty

Neurology 2006;67;E19

DOI 10.1212/01.wnl.0000244494.50684.5b

This information is current as of November 27, 2006

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/67/10/E19.full
References	This article cites 1 articles, 1 of which you can access for free at: http://n.neurology.org/content/67/10/E19.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Cerebrovascular disease/Stroke http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke DWI http://n.neurology.org/cgi/collection/dwi Infarction http://n.neurology.org/cgi/collection/infarction MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

