


Teaching NeuroImages: Primary diffuse large B-cell lymphoma of the cranial vault

D. Renard, MD
C. Campello, MD
O. Beraru, MD
P. Bouillot, MD
P. Labauge, MD, PhD

Address correspondence and
reprint requests to Dr. Dimitri
Renard, Department of
Neurology, CHU Nîmes, Hôpital
Caremeau, Place du Pr Debré,
30029 Nîmes Cedex 4, France
dimitrirenard@hotmail.com

Figure Brain images showing an intracranial and extracranial frontal mass lesion associated with local cranial bone destruction



Brain CT showing a slightly hyperintense intracranial and extracranial frontal mass lesion (A) with homogeneous enhancement after contrast injection (B), associated with local cranial bone destruction (C). This lesion was isointense on T1-weighted (D) and T2-weighted (E) MRI. Gadolinium-enhanced T1 sequences showed strong, slightly heterogeneous enhancement (F).

A 67-year-old woman presented with a painful right-sided scalp swelling. Brain imaging showed an intracranial and extracranial frontal mass lesion associated with local cranial bone destruction (figure). The radiologic differential diagnosis included malignant meningioma, primary bone tumor, and bone metastasis. Thoracic, abdominal, pelvic, and orbital CT scan, bone scintigraphy, whole-body fluorodeoxyglucose PET, bone marrow biopsy, lumbar puncture, and ophthalmologic

examination (including slit lamp examination) showed no other lesions. Open lesion biopsy revealed a diffuse malignant large B-cell non-Hodgkin lymphoma of the dura mater and cranial bone. Six cycles of R-CHOP (rituximab with cyclophosphamide, doxorubicin, vincristine, and prednisone) regimen was started. Primary bone (most frequently non-Hodgkin) lymphoma are rare.^{1,2} In our case, associated dura mater involvement (which seemed to be secondary)

From the Department of Neurology (D.R., C.C., O.B., P.L.), CHU Nîmes, Hôpital Caremeau; and Department of Neurosurgery (P.B.), Clinique des Franciscaines, Nîmes, France.

Disclosure: The authors report no disclosures.

was seen on both brain imaging and histology. Primary bone tumors often have a favorable outcome, especially when treated by combined modality therapy. Clinical stage is the most important prognostic variable in predicting overall survival.

REFERENCES

1. Aquilina K, O'Brian DF, Phillips JP. Diffuse primary non-Hodgkin's lymphoma of the cranial vault. *Br J Neurosurg* 2004;18:518–523.
2. Agbi CB, Bannister CM, Turnbull IW. Primary cranial vault lymphoma mimicking a meningioma. *Neurochirurgia* 1983;26:130–132.

Neurology[®]

Teaching *NeuroImages*: Primary diffuse large B-cell lymphoma of the cranial vault

D. Renard, C. Campello, O. Beraru, et al.

Neurology 2009;73:e84-e85

DOI 10.1212/WNL.0b013e3181bd8283

This information is current as of October 26, 2009

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/73/17/e84.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/73/17/e84.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Oncology http://n.neurology.org/cgi/collection/all_oncology CT http://n.neurology.org/cgi/collection/ct MRI http://n.neurology.org/cgi/collection/mri Primary brain tumor http://n.neurology.org/cgi/collection/primary_brain_tumor
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

