RESIDENT & FELLOW SECTION

Section Editor Mitchell S.V. Elkind, MD, MS

Gayatri S. Reilly, MD Robert K. Shin, MD

Address correspondence and reprint requests to Dr. Robert K. Shin, Department of Neurology, University of Maryland, 110 S. Paca Street, Third Floor, Baltimore, MD 21201 rshin@som.umaryland.edu

Teaching Neuro *Images*: Herpes zoster ophthalmicus—related oculomotor palsy accompanied by Hutchinson sign

Figure 1

Partial ptosis, mydriasis, and exotropia consistent with a right oculomotor palsy



The skin lesion on the tip of the nose (Hutchinson sign) signifies involvement of the nasociliary branch of V1, which also innervates ocular structures.

A 51-year-old woman presented with acute diplopia. Findings include right ptosis, a dilated, unreactive pupil, and impaired adduction and vertical ductions (figure 1). A skin lesion was noted on the right tip of the nose, residual from a vesicular rash over the right forehead 3 weeks earlier (figure 1). MRI demonstrated enhancement of the cisternal third nerve, obviating the need for angiography (figure 2). The oculomotor palsy resolved within 3 months.

Oculomotor palsy may present weeks after herpes zoster ophthalmicus.^{1,2} Involvement of the tip of the nose (Hutchinson sign) is a strong predictor of ocular involvement, indicating involvement of the nasociliary branch of V1, which innervates both the tip of the nose and ocular structures such as the cornea, conjunctiva, and uvea.

Figure 2

T1-weighted MRI showing enhancement of the cisternal portion of the oculomotor nerve (arrow)



REFERENCES

- Chang-Godinich A, Lee AG, Brazis PW, Liesegang TJ, Jones DB. Complete ophthalmoplegia after zoster ophthalmicus. J Neuroophthalmol 1997;17:262–265.
- Quisling SV, Shah VA, Lee HK, et al. Magnetic resonance imaging of third cranial nerve palsy and trigeminal sensory loss caused by herpes zoster. J Neuroophthalmol 2006;26: 47–48.



Teaching Neuro Images: Herpes zoster ophthalmicus—related oculomotor palsy accompanied by Hutchinson sign

Gayatri S. Reilly and Robert K. Shin *Neurology* 2010;74;e65 DOI 10.1212/WNL.0b013e3181d8c1f6

This information is current as of April 12, 2010

Updated Information & including high resolution figures, can be found at:

Services http://n.neurology.org/content/74/15/e65.full

References This article cites 2 articles, 0 of which you can access for free at:

http://n.neurology.org/content/74/15/e65.full#ref-list-1

Subspecialty Collections This article, along with others on similar topics, appears in the

following collection(s):
All Neuro-ophthalmology

http://n.neurology.org/cgi/collection/all_neuroophthalmology

Diplopia (double vision)

http://n.neurology.org/cgi/collection/diplopia_double_vision

Ocular motility

http://n.neurology.org/cgi/collection/ocular_motility

Post-infectious

http://n.neurology.org/cgi/collection/postinfectious_

Pupils

http://n.neurology.org/cgi/collection/pupils

Permissions & Licensing Information about reproducing this article in parts (figures, tables) or in

its entirety can be found online at:

http://www.neurology.org/about/about_the_journal#permissions

Reprints Information about ordering reprints can be found online:

http://n.neurology.org/subscribers/advertise

Neurology ® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright . All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

