

fracture risk assessment (FRAX) and falls assessment annually, with a baseline bone mineral density scan to identify those at high risk of osteoporotic fracture. Potential contributors to fracture risk should be avoided where possible, and interventions to improve both bone health and falls risk should be routine.

Author Response: Marloes T. Bazelier, Frank de Vries, Utrecht, the Netherlands: We appreciate the comments by Dobson et al. and agree with their concerns about the use of anxiolytics/hypnotics and antidepressants in patients with MS. These medication types have been associated with falls and (hip) fractures.^{7,8} However, there is no evidence that discontinuation of these drugs would prevent fractures. We also agree that epidemiologic evidence for the underlying etiology of glucocorticoid use and risk of fractures in patients with MS is unclear.^{4,6} Because patients with MS are already at risk of fracture, FRAX scores may be underestimated. Unfortunately, FRAX has not been designed specifically for patients with MS. We have recently published a clinical risk score that has been developed for fracture risk assessment in patients with MS.⁹

© 2012 American Academy of Neurology

1. Bazelier MT, van Staa TP, Uitdehaag BM, et al. Risk of fractures in patients with multiple sclerosis: a population-based cohort study. *Neurology* 2012;78:1967–1973.
2. Ensrud KE, Blackwell TL, Mangione CM, et al. Central nervous system-active medications and risk for falls in older women. *J Am Geriatr Soc* 2002;50:1629–1637.
3. Vestergaard P, Rejnmark L, Mosekilde L. Anxiolytics, sedatives, antidepressants, neuroleptics and the risk of fracture. *Osteoporos Int* 2006;17:807–816.
4. Bazelier MT, van Staa T, Uitdehaag BM, et al. The risk of fracture in patients with multiple sclerosis: the UK general practice research database. *J Bone Miner Res* 2011;26:2271–2279.
5. Schwid SR, Goodman AD, Puzas JE, McDermott MP, Mattson DH. Sporadic corticosteroid pulses and osteoporosis in multiple sclerosis. *Arch Neurol* 1996;53:753–757.
6. Bazelier MT, Bentzen J, Vestergaard P, et al. The risk of fracture in incident multiple sclerosis patients: The Danish National Health Registers. *Mult Scler* 2012 Epub Apr 3.
7. Verdel BM, Souverein PC, Egberts TC, van Staa TP, Leufkens HG, de Vries F. Use of antidepressant drugs and risk of osteoporotic and non-osteoporotic fractures. *Bone* 2010;47:604–609.
8. van den Brand MW, Pouwels S, Samson MM, et al. Use of antidepressants and the risk of fracture of the hip or femur. *Osteoporos Int* 2009;20:1705–1713.
9. Bazelier MT, van Staa TP, Uitdehaag BM, et al. A simple score for estimating the long-term risk of fracture in patients with multiple sclerosis. *Neurology* 2012;79:922–928.

CORRECTIONS

Plasma multianalyte profiling in mild cognitive impairment and Alzheimer disease

In the article “Plasma multianalyte profiling in mild cognitive impairment and Alzheimer disease” by W.T. Hu et al. (*Neurology*® 2012;79:897–905), there is an error in the first paragraph on page 899. The third sentence should read “At WU, blood was collected in EDTA in polypropylene tubes after overnight fasting between 7:30 and 8:00 AM and centrifuged (2,000 g × 15 minutes at 4°C) for separation into plasma and cellular components.” The authors regret the error.

WriteClick: Editor’s Choice: Predicting outcome after acute basilar artery occlusion based on admission characteristics

In the correspondence regarding the article “Predicting outcome after acute basilar artery occlusion based on admission characteristics” by Y. He et al. (*Neurology*® 2012;79:1410), there is an error in the second author’s name, which should be spelled “Tianxiao Li.” The editorial staff regrets the error.

Author disclosures are available upon request (journal@neurology.org).

Neurology[®]

**WriteClick: Editor's Choice: Predicting outcome after acute basilar artery occlusion
based on admission characteristics**

Neurology 2012;79;1935
DOI 10.1212/WNL.0b013e3182771ed7

This information is current as of October 29, 2012

**Updated Information &
Services**

including high resolution figures, can be found at:
<http://n.neurology.org/content/79/18/1935.2.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in
its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2012 by AAN Enterprises, Inc.. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

